

An allosteric model of calmodulin explains differential activation of PP2B and CaMKII

Melanie I. Stefan, Stuart J. Edelstein, and Nicolas Le Novère*

European Molecular Biology Laboratory–European Bioinformatics Institute,
Wellcome Trust Genome Campus, Hinxton CB10 1SD, United Kingdom

The version of the article published by the *Proceedings of the National Academy of Science USA* contains ancient versions of Figure 3 and Figure 4, obtained with an imperfect parameter search approach. The substitution does not change the scientific message of the article. However, the published figures are far less convincing than the correct ones. The correct Figure 3 shows a better fit between our model and experimental data, while the correct Figure 4 demonstrates a far larger effect of the target on Calmodulin affinity for calcium. Correct versions are appended to the present document, after the reprint from the *PNAS*.

An allosteric model of calmodulin explains differential activation of PP2B and CaMKII

Melanie I. Stefan, Stuart J. Edelstein, and Nicolas Le Novère*

European Molecular Biology Laboratory–European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton CB10 1SD, United Kingdom

Communicated by Jean-Pierre Changeux, Institut Pasteur, Paris, France, May 15, 2008 (received for review January 15, 2008)

Calmodulin plays a vital role in mediating bidirectional synaptic plasticity by activating either calcium/calmodulin-dependent protein kinase II (CaMKII) or protein phosphatase 2B (PP2B) at different calcium concentrations. We propose an allosteric model for calmodulin activation, in which binding to calcium facilitates the transition between a low-affinity [tense (*T*)] and a high-affinity [relaxed (*R*)] state. The four calcium-binding sites are assumed to be nonidentical. The model is consistent with previously reported experimental data for calcium binding to calmodulin. It also accounts for known properties of calmodulin that have been difficult to model so far, including the activity of nonsaturated forms of calmodulin (we predict the existence of open conformations in the absence of calcium), an increase in calcium affinity once calmodulin is bound to a target, and the differential activation of CaMKII and PP2B depending on calcium concentration.

allostery | synaptic plasticity | calcium binding | cooperativity | conformational transition

Activity-dependent changes in synaptic strength (1) have long been used as a paradigm to study learning and memory (reviewed in ref. 2). Calcium signaling is a key factor in both long-lasting increases [known as long-term potentiation (LTP)] and long-lasting decreases [long-term depression (LTD)] in synaptic strength. According to a model first proposed by Lisman (3), the coordinated activity of a pair of neurons leads to a large increase in calcium levels in the postsynaptic neuron and an increase in synaptic strength, whereas the activity of only one of the two neurons results in more moderate postsynaptic calcium levels and, consequently, a reduction in synaptic strength. Calcium entering the postsynaptic neuron through NMDA receptors and voltage-operated calcium channels and from the endoplasmic reticulum activates calmodulin. Activated calmodulin may bind to calcium/calmodulin kinase II (CaMKII) and increases its activity (4–6). Active CaMKII enhances the function of AMPA receptor channels by phosphorylating the GluR1 subunit (7). It also mediates an increase of AMPA receptor delivery to the postsynaptic membrane (8). These roles are consistent with reports implicating CaMKII in some forms of learning and memory (9). In contrast, lower amounts of calcium in the postsynaptic neuron will cause calmodulin to activate PP2B, leading to activation of protein phosphatase 1 and a subsequent reduction of CaMKII activity (reviewed in ref. 10). It remains to be explained, however, how calmodulin performs this dual function dependent on calcium levels.

Calmodulin is a ubiquitous regulatory protein that binds four calcium ions (11, 12). It is a single polypeptide chain of 148-aa residues (13) and can adopt two distinct conformations: in the absence of calcium, its EF hands typically adopt an inactive, compact (closed) form (14). When bound to four calcium ions, they are found in an open active form (15).

A variety of models for calmodulin activation and action have been used in the past. Each of these models reflects some properties of calmodulin and is reasonably applicable in contexts in which only these properties are relevant. However, none of these models can satisfactorily account for all of the observed properties of calmodulin such as cooperativity of calcium bind-

ing and different affinities for different calcium-binding sites (16), activation of targets by unsaturated calmodulin (17, 18), and increased affinity for calcium upon binding to targets (18–20). We propose an alternative model, based on a biophysical description of the conformational transitions. Originally applied to oligomeric proteins with symmetric identical subunits (21), this approach can also be adapted to a single polypeptide chain with multiple binding sites. The resulting generalized allosteric model of calmodulin can reconcile different properties of calmodulin, including differential activation of PP2B and CaMKII, residual activation of CaMKII at low calcium concentration, differences between the binding sites in terms of calcium affinity, and the existence of active and inactive conformations.

Allosteric Model of Calmodulin. In our model, calmodulin can exist in two different states, the active open [relaxed (*R*)] state and the inactive closed [tense (*T*)] state. Each of these states can bind four calcium ions (Fig. 1). When no calcium is bound, the *T* state prevails, because its free energy is lower than that of the unbound *R* state. Consecutive binding of calcium ions, however, progressively stabilizes the *R* state until the free energy of the *R* state is lower than that of the *T* state, so the *R* state is favored.

The four different binding sites are designated *A*, *B*, *C*, and *D* (*A* and *B* on the N-terminal domain, *C* and *D* on the C-terminal domain, with no sequential order being implied within the domains). Each of the states and each of the reactions is explicitly modeled, with distinct dissociation constants and *R* to *T* transition probabilities for each of the sites (Fig. 2). The constant *L* describes the equilibrium between both states when no calcium ion is bound: $L = [T_0]/[R_0]$. If *L* is very large, most of the protein exists in the tense state in the absence of calcium. If *L* is small (close to one), the *R* state is nearly as populated as the *T* state. The constants c_A , c_B , c_C , and c_D describe the ratio of dissociation constants for the *R* and *T* states for each site: $c_i = K_i^R/K_i^T$. If *c* is one, both *R* and *T* states have the same affinity for calcium. The *c* values also indicate how much the equilibrium between *T* and *R* states changes upon calcium binding: the smaller *c*, the more the equilibrium shifts toward the *R* state.

The formula for fractional occupation of an allosteric protein in the absence of allosteric effectors (21) can be generalized to describe nonequivalent calcium-binding sites. In the case of four binding sites, the generalized expression is:

$$\bar{Y} = 0.25 \frac{\sum_i \left(\alpha_i \prod_j (1 + \alpha_j) \right) + L \sum_i \left(c_i \alpha_i \prod_j (1 + c_j \alpha_j) \right)}{\prod_i (1 + \alpha_i) + L \prod_i (1 + c_i \alpha_i)}, \quad [1]$$

Author contributions: M.I.S. and N.L. designed research; M.I.S. and N.L. performed research; M.I.S. and S.J.E. analyzed data; and M.I.S., S.J.E., and N.L. wrote the paper.

The authors declare no conflict of interest.

Data deposition: The model reported in this paper has been deposited in the BioModels database, www.ebi.ac.uk/biomodels (accession code MODEL9885984404).

*To whom correspondence should be addressed. E-mail: lnov@ebi.ac.uk.

This article contains supporting information online at www.pnas.org/cgi/content/full/0804672105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA

the fact that targets act as allosteric activators, drawing the equilibrium toward the high-affinity *R* state.

Note that a simple model based on thermal equilibrium has been proposed before (50), although based on sequential bindings of calcium. Furthermore, the author did not try to estimate parameters using experimental information or to validate the model.

Model Characteristics. The association constants we determined from the simulation result by fitting it with an Adair-type equation (43) are the observable sequential association constants for the first, second, third, and fourth binding events. It is important to note, however, that our model does not assume a fixed order for calcium binding to the different binding sites. Rather, because the first calcium ion can bind to any one of four binding sites in either state, K_1 is a combination of the microscopic K_i^R and K_i^T values for each of the sites in each of the states. The apparent sequential dissociation constants used in our model follow an Adair-type (43) framework as used, for instance, by Crouch and Klee (16) and Porumb (30). They are, in principle, Adair constants, except for a slight difference in nomenclature: K_2 according to Adair (43) corresponds to $K_1 \times K_2$ according to Crouch and Klee (16) and Porumb (30), and so on.

It is important to note that the conformation of calmodulin which we call the *T* state is not necessarily exactly identical to the reported *apo* structure (14) of calmodulin. Rather, the *T* state represents a collection of structures that may differ somewhat in the conformation of the calcium-binding sites, but whose overall structure resembles that of *apo* calmodulin. The existence of an ion-bound form that resembles the *apo* conformation has recently been established (51). Likewise, the *R* state is a collection of structures that resemble the reported open structure of active calmodulin (52, 53). Asymmetric forms of calmodulin with one lobe in an open state and one head in a closed state have been

reported in the presence of some targets (54). However, the binding mechanism of these targets differs from that of CaMKII and PP2B, where both lobes bind to the target.


Lisman Hypothesis. The model proposed here explains how different amounts of calcium can trigger the activation of PP2B or CaMKII and thus provides support for the Lisman hypothesis (3) on a molecular level. The question of how different frequencies of calcium signals lead to differential activation of PP2B or CaMKII is not addressed in the model. It has been suggested, however, that, at least under some conditions, high frequencies of calcium input result in high local concentrations of calcium, whereas low calcium frequencies result in moderate local calcium concentrations in the spine (44, 55, 56). In addition, calcium frequency has a direct impact on CaMKII, because of the requirement for two adjacent subunits to be active for auto-phosphorylation at threonine residue 286, which confers sustained activity (57–59). Other factors the model does not account for include variations in the subcellular localization of PP2B (reviewed in ref. 60) and CaMKII (61) and the inhibitory effect of PP2B on CaMKII (reviewed in ref. 10). The latter effect, if included, would increase the window of calcium concentrations at which PP2B is preferably activated, enhancing the distinction between PP2B and CaMKII activation. We believe, however, that our model provides a valid and useful biophysical basis on which to develop further models of synaptic plasticity mechanisms.

Note added in proof (64). Experimental support for our equilibrium model of calmodulin function has been recently published by Gsponer *et al.*

ACKNOWLEDGMENTS. We thank Julia Shifman (Hebrew University of Jerusalem) for sharing raw data on calcium binding to calmodulin in the presence and absence of CaMKII. We also thank Laurence Calzone, Stephen Martin, and Annalisa Pastore for helpful comments on the manuscript. S.J.E. was supported by an International Short Visit Fellowship from the Royal Society.

- Bliss TV, Lomo T (1973) Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol* 232:331–356.
- Martin SJ, Grimwood PD, Morris RG (2000) Synaptic plasticity and memory: An evaluation of the hypothesis. *Annu Rev Neurosci* 23:649–711.
- Lisman J (1989) A mechanism for the Hebb and the anti-Hebb processes underlying learning and memory. *Proc Natl Acad Sci USA* 86:9574–9578.
- Schulman H, Greengard P (1978) Stimulation of brain membrane protein phosphorylation by calcium and an endogenous heat-stable protein. *Nature* 271:478–479.
- Schulman H, Greengard P (1978) Ca²⁺-dependent protein phosphorylation system in membranes from various tissues, and its activation by "calcium-dependent regulator." *Proc Natl Acad Sci USA* 75:5432–5436.
- Lisman J, Schulman H, Cline H (2002) The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat Rev Neurosci* 3:175–190.
- Lee HK, Barbarosie M, Kameyama K, Bear MF, Huganir RL (2000) Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* 405:955–959.
- Hayashi Y, *et al.* (2000) Driving AMPA receptors into synapses by LTP and CaMKII: Requirement for GluR1 and PDZ domain interaction. *Science* 287:2262–2267.
- Silva AJ, Paylor R, Wehner JM, Tonegawa S (1992) Impaired spatial learning in alpha-calmodulin-calmodulin kinase II mutant mice. *Science* 257:206–211.
- Groth RD, Dunbar RL, Mermelstein PG (2003) Calcineurin regulation of neuronal plasticity. *Biochem Biophys Res Commun* 311:1159–1171.
- Lin YM, Liu YP, Cheung WY (1974) Cyclic 3':5'-nucleotide phosphodiesterase. Purification, characterization, and active form of the protein activator from bovine brain. *J Biol Chem* 249:4943–4954.
- Xia Z, Storm DR (2005) The role of calmodulin as a signal integrator for synaptic plasticity. *Nat Rev Neurosci* 6:267–276.
- Watterson DM, Sharief F, Vanaman TC (1980) The complete amino acid sequence of the Ca²⁺-dependent modulator protein (calmodulin) of bovine brain. *J Biol Chem* 255:962–975.
- Kuboniwa H, *et al.* (1995) Solution structure of calcium-free calmodulin. *Nat Struct Biol* 2:768–776.
- Babu YS, *et al.* (1985) Three-dimensional structure of calmodulin. *Nature* 315:37–40.
- Crouch TH, Klee CB (1980) Positive cooperative binding of calcium to bovine brain calmodulin. *Biochemistry* 19:3692–3698.
- Kincaid RL, Vaughan M (1986) Direct comparison of Ca²⁺ requirements for calmodulin interaction with and activation of protein phosphatase. *Proc Natl Acad Sci USA* 83:1193–1197.
- Shifman JM, Choi MH, Mihalas S, Mayo SL, Kennedy MB (2006) Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) is activated by calmodulin with two bound calciums. *Proc Natl Acad Sci USA* 103:13968–13973.
- Burger D, Stein EA, Cox JA (1983) Free energy coupling in the interactions between Ca²⁺, calmodulin, and phosphorylase kinase. *J Biol Chem* 258:14733–14739.
- Olwin BB, Edelman AM, Krebs EG, Storm DR (1984) Quantitation of energy coupling between Ca²⁺, calmodulin, skeletal muscle myosin light chain kinase, and kinase substrates. *J Biol Chem* 259:10949–10955.
- Monod J, Wyman J, Changeux JP (1965) On the nature of allosteric transitions: A plausible model. *J Mol Biol* 12:88–118.
- Meador WE, Means AR, Quirocho FA (1992) Target enzyme recognition by calmodulin: 2.4 Å structure of a calmodulin-peptide complex. *Science* 257:1251–1255.
- Ye Q, Li X, Wong A, Wei Q, Jia Z (2006) Structure of calmodulin bound to a calcineurin peptide: A new way of making an old binding mode. *Biochemistry* 45:738–745.
- Rubin MM, Changeux JP (1966) On the nature of allosteric transitions: Implications of non-exclusive ligand binding. *J Mol Biol* 21:265–274.
- Peersen OB, Madsen TS, Falke JJ (1997) Intermolecular tuning of calmodulin by target peptides and proteins: Differential effects on Ca²⁺ binding and implications for kinase activation. *Protein Sci* 6:794–807.
- Bailey PM, Findlay WA, Martin SR (1996) Target recognition by calmodulin: Dissecting the kinetics and affinity of interaction using short peptide sequences. *Protein Sci* 5:1215–1228.
- Hoops S, *et al.* (2006) COPASI—a COmplex PAthway Simulator. *Bioinformatics* 22:3067–3074.
- Burger D, Cox JA, Comte M, Stein EA (1984) Conformational changes in calmodulin upon binding of calcium. *Biochemistry* 23:1966–1971.
- Haiech J, Klee CB, Demaille JG (1981) Effects of cations on affinity of calmodulin for calcium: Ordered binding of calcium ions allows the specific activation of calmodulin-stimulated enzymes. *Biochemistry* 20:3890–3897.
- Porumb T (1994) Determination of calcium-binding constants by flow dialysis. *Anal Biochem* 220:227–237.
- Quintana AR, Wang D, Forbes JE, Waxham MN (2005) Kinetics of calmodulin binding to calcineurin. *Biochem Biophys Res Commun* 334:674–680.
- Tzortzopoulos A, Török K (2004) Mechanism of the T286A-mutant alphaCaMKII interactions with Ca²⁺/calmodulin and ATP. *Biochemistry* 43:6404–6414.
- Collins MO, *et al.* (2006) Molecular characterization and comparison of the components and multiprotein complexes in the postsynaptic proteome. *J Neurochem* 97:16–23.
- Chen X, *et al.* (2005) Mass of the postsynaptic density and enumeration of three key molecules. *Proc Natl Acad Sci USA* 102:11551–11556.

Correct figure 3



Correct figure 4

