




MEDICAL HOME

LIFE SCIENCES HOME

Become a Member

Search...

About News Health A-Z Drugs Medical Devices Interviews White Papers MediKnowledge More...



Our brain senses angry voices in a flash, study shows

Download PDF Copy

Reviewed by *Alina Shrourou, BSc*

Dec 7 2018

Sight and hearing are the two main sensory modalities allowing us to interact with our environment. But what happens within the brain when it perceives a threatening signal, such as an aggressive voice? How does it distinguish a threatening voice from the surrounding noise? How does it process this information? To answers these questions, researchers from the University of Geneva (UNIGE), Switzerland, studied brain activity during the processing of various emotional voices. They discovered that we notice a voice much faster when it is considered threatening than when it is perceived as normal or happy. Our attention is more focused on threatening voices to enable us to clearly recognize the location of the potential threat. This study, published in the journal *Social, Cognitive and Affective Neuroscience*, demonstrates the resources leveraged by our brain when we sense danger to allow for adequate survival behavior.

Sight and hearing are the two senses that allow human beings to detect threatening situations. Although sight is critical, it does not allow for a 360-degree coverage of the surrounding space - unlike hearing. "That's why we are interested in how fast our attention responds to the different intonations of the voices around us and how our brain deals with potentially threatening situations," explains Nicolas Burra, a researcher in the psychology section of the Faculty of Psychology and Education Sciences (FPSE) at UNIGE.

To examine the brain's response to threats in the auditory environment, the researchers presented 22 short human voice sounds (600 milliseconds) that were neutral utterances or expressed either anger or joy. Using two loudspeakers, these sounds were presented to 35 participants while an electroencephalogram (EEG) measured electrical activity in the brain down to the millisecond. More specifically, the researchers focused on the electrophysiological components related to auditory attentional processing. "Each participant heard two sounds simultaneously: two neutral voices, one neutral and one angry voice, or one

Advertisement

Presents Andrea Oliva

Am Donnerstag 13. 12. startet das grosse dreitägige Pop Up-Restaurant-Konzept Onagi

Onagi Restaurant Clubbing

- Trending Stories
- Latest Interviews
- Top Health Articles

Gene that allows you eat as much as you want could help combat obesity

From Our Bench to Yours: Antibody Advice from Proteintech

LetsGetChecked: The Company That's Revolutionizing Consumer Healthcare

Researchers explore why and how Mediterranean diet may mitigate cardiovascular risk

Brown researchers awarded \$12 million NIH grant renewal to explore fundamentals of brain function

[Lire en ligne](#)

neutral and one happy voice. When they perceived anger or joy, they had to respond by pressing a key on a keyboard as accurately and quickly as possible," explains Leonardo Ceravolo, researcher at UNIGE's Swiss Centre for Affective Sciences. "We then measured the intensity of brain activity when attention is focused on the different sounds, as well as the duration of this focus before a return to the basic state," he adds.

Our brain quickly differentiates angry from happy voices

Using data from the EEG, the researchers examined the appearance of a cerebral marker of auditory attention called N2ac. As Nicolas Burra explains, "When the brain perceives an emotional target sound, N2ac activity is triggered after 200 milliseconds. However, when it perceives anger, the N2ac is amplified and lasts longer, which is not the case for joy!"

Related Stories

- Mapping brain microglia
- Maryland researchers receive grant to study the brain-to-gut connection in schizophrenia
- Combining toxic herbicide with sugar-binding proteins may trigger Parkinsonism

Subsequently, after 400 milliseconds, our attention must disengage from the emotional vocal stimulus. At this moment, a cerebral marker of auditory attention, called LPCpc, intervenes. Interestingly enough, LPCpc activity is also stronger for angry than for happy voices. Why? "Anger can signal a potential threat, which is why the brain analyzes these kinds of stimuli for a longer time. In an auditory environment, this mechanism allows us to not become alarmed at the slightest potentially threatening noise or, conversely, to adopt the most appropriate behavior in case of danger. These extra milliseconds of attention are, therefore, crucial to the accurate interpretation of a threat in a complex auditory environment," says Ceravolo.

This additional temporal cost was also evident in the participants' response times. When they had to indicate that they perceived anger, it took them longer than when they did so for joy. In contrast, brain activity was enhanced in the case of angry stimuli. Does it sound conflicting? "No. The explanation is logical. As attention in the brain remains focused on the threatening sound, the motor response via the keyboard is delayed," says Nicolas Burra.

In summary, this study demonstrated for the first time that in a few hundred milliseconds, our brain is sensitive to the presence of angry voices. This rapid

Neu: Onagi Restaurant Clubbing

Onagi Restaurant Clubbing

Erstmal in der Schweiz wird ein Burger mit japanischem Wagyu A5+ Fleisch serviert.

ÖFFNEN

Advertisement

Latest News

HPV type 16 or 18 associated with cervical cancer risk in young women

New artificial intelligence-based system to differentiate between different types of cancer cells

Cervical cancer risk is higher in women with positive HPV, but no cellular abnormalities



detection of the source of a potential threat in a complex environment is essential, as it is "critical in crisis situations and a great advantage for our survival," concludes Ceravolo.

Study compares pain-related diagnoses in First Nations and non-First Nations children, youth

Source:

https://www.unige.ch/

Updated breast cancer screening guideline highlights importance of shared decision-making

Be the first to rate this article



Posted in: [Medical Science News](#) | [Medical Research News](#)

Tags: [Brain](#), [Education](#), [Neuroscience](#), [Psychology](#)

Comments (0)

Download PDF Copy

Try the Maxwell® RSC PureFood and GMO Authentication Kit for Purifying DNA for Food and Ingredient Authentication



Advertisement

Read in: [English](#)

News Medical 268,550 likes
Like Page Share
Be the first of your friends to like this

Suggested Reading

Newsletters you may be interested in



Neurology / Neuroscience
(Subscribe or Preview)



Psychiatry
(Subscribe or Preview)



Sensors
(Subscribe or Preview)

See all Newsletters »

FinnBrain imaging study explores brain structure in newborns

Novel approach gives new insights into formation of neurons in mammalian adult brain

Study reveals possible link between hysterectomy and brain function

New EU-funded project provides insight into how the brain develops

Cedars-Sinai research identifies neurons that play role in how people recognize errors

Researchers find effective new target for mood-boosting brain stimulation

Study reveals why some children are susceptible to the effects of maltreatment

Brown researchers awarded \$12 million NIH grant renewal to explore fundamentals of brain function

Comments



[Lire en ligne](#)

The opinions expressed here are the views of the writer and do not necessarily reflect the views and opinions of News-Medical.Net.



Post a new comment

Login

Quirky Comment Title (optional)

Post

Medical Links

Medical Home
News
Health A-Z
White Papers
Thought Leaders
Insights

MediKnowledge Series
Health & Personal Care
Medical Devices
Drugs

Life Sciences Links

Life Sciences Home
News
Lab Instruments & Equipment
Life Sciences A-Z
White Papers
Thought Leaders
Insights
Webinars

Other Useful Links

About
Meet the Team
Search
Newsletters
Twitter Channels
Sitemap
Advertise
Contact
Update Your Privacy Preferences
Terms & Conditions
Privacy & Cookie Policy



News-Medical.Net provides this medical information service in accordance with these [terms and conditions](#). Please note that medical information found on this website is designed to support, not to replace the relationship between patient and physician/doctor and the medical advice they may provide.



512 people on this site
powered by chartbeat

This site complies with the HONcode standard for trustworthy health information: [verify here](#).



By continuing to browse or by clicking "Accept All Cookies," you agree to the storing of first and third-party cookies on your device to enhance site navigation, analyze site usage, and assist in our marketing efforts. [Find out more](#).

[Cookie Settings](#)

Accept All Cookies

