




under embargo until October 26th, 7pm, Swiss time

RESEARCHERS DECIPHER THE MECHANISM OF MEMBRANE FISSION

Thanks to a detailed analysis of the role of dynamin, a team of researchers explains the phenomenon of how membranes break



Artistic view of a dynamin helix around an endocytosis bud

Credit: Nymus3D

Researchers at the University of Geneva (UNIGE), in collaboration with the Institut Curie (Paris, France), shed light on the mechanism of the fission of cell membranes. They used an in vitro analysis of the functioning of dynamin, a protein involved in this mechanism. The results were published in the journal *Cell*.

A cell is composed of a nucleus which encloses its genetic information and the cytoplasm which is itself confined by an external membrane separating the cell from the outside world. The impermeability of the membrane and its ability to repair itself protect the cell from its environment. Although this membrane resistance is fundamental to the survival of the cell, the cell also needs to let in particles necessary for its proper functioning. The mechanism by which a small region of the cytoplasmic membrane invaginates to form a bud that will then be sectioned off to let molecules and other particles into the cell is known as endocytosis.

However, this natural process remains elusive due to the remarkable resistance of the cell membrane. Aurélien Roux, a professor of biochemistry and member of the National Centres of Competence in Research (NCCR) Chemical Biology, heads a team that focused on dynamin, a protein involved in endocytosis, to try to understand how an ultra-resistant membrane can nevertheless let external elements enter into the cell.

The power of dynamin

Scientists conducted in vitro experiments using artificial membrane tubules with a radius of 10 to 100 nanometres. They discovered that once dynamin is injected into the tube, it polymerises. In other words, it forms a helix around the tube and compresses it until it breaks. Dynamin produces the energy necessary for this constriction by «consuming» GTP molecules, much like a car consumes gasoline.

Based on these experiments, Professor Roux's team observed that the location of the fission is very specific and appears at the boundary between the helix and the membrane. «A change in radius that curves the membrane, caused by the polymerisation of dynamin, induces a stress that promotes the fracture,» states Sandrine Morlot, researcher at the Department of Biochemistry. «This is new data allowing us to explain the process of fission.»

The researchers were also able to measure the time it took to fission the membrane. Its duration depends on the mechanical properties of the membrane, which vary from one cell to another.

the «**dynamine**» is a protein involved in the endocytosis phenomenon

«We found that the ability of dynamine to break an ultra-resistant membrane is due to its torque, that is to say, its rotational force, which is vastly superior to that of other proteins,» explains Professor Roux. «By decrypting the effect of dynamine on the membrane, we have come to understand the workings of membrane fission, a phenomenon which is certainly natural but remains extremely complex.»

contact

Aurélien Roux

022 379 35 32

aurelien.roux@unige.ch

UNIVERSITÉ DE GENÈVE
Service de communication

24 rue du Général-Dufour
CH-1211 Genève 4

Tél. 022 379 77 17

media@unige.ch

www.unige.ch