

A microscopic derivation of time-dependent correlation functions of the 1D cubic nonlinear Schrödinger equation

Jürg Fröhlich* Antti Knowles† Benjamin Schlein‡ Vedran Sohinger§

January 26, 2019

We give a microscopic derivation of time-dependent correlation functions of the 1D cubic nonlinear Schrödinger equation (NLS) from many-body quantum theory. The starting point of our proof is [11] on the time-independent problem and [16] on the corresponding problem on a finite lattice. An important new obstacle in our analysis is the need to work with a cutoff in the number of particles, which breaks the Gaussian structure of the free quantum field and prevents the use of the Wick theorem. We overcome it by means of complex analytic methods. Our methods apply to the nonlocal NLS with bounded convolution potential. In the periodic setting, we also consider the local NLS, arising from short-range interactions in the many-body setting. To that end, we need the dispersion of the NLS in the form of periodic Strichartz estimates in $X^{s,b}$ spaces.

1. Setup and main result

Let \mathfrak{H} be a Hilbert space, $H \in C^\infty(\mathfrak{H})$ a Hamiltonian function, and $\{\cdot, \cdot\}$ a Poisson bracket on $C^\infty(\mathfrak{H}) \times C^\infty(\mathfrak{H})$. We can then define the Hamiltonian flow of H on \mathfrak{H} , which we denote by $u \mapsto S_t u$. Furthermore, we introduce the *Gibbs measure* associated with the Hamiltonian H , defined as a probability measure \mathbb{P} on \mathfrak{H} formally given by

$$d\mathbb{P}(u) := \frac{1}{Z} e^{-H(u)} du, \quad (1.1)$$

where Z is a positive normalization constant and du is Lebesgue measure on \mathfrak{H} (which is ill-defined if \mathfrak{H} is infinite-dimensional). The problem of the construction of measures of the type (1.1) was first considered in the constructive quantum field theory literature, c.f. [13, 24] and the references therein, and later in [17, 21, 22]. In the context of nonlinear dispersive PDEs, the invariance of measures of the type (1.1) has been considered in the work of Bourgain [4–8] and Zhidkov [28], and in the subsequent literature. An important application of the invariance is to obtain a substitute for a conservation law at low regularity which, in turn, allows us to construct solutions for random initial data of low regularity. We refer the reader to the introduction of [11] for a detailed overview and for further references.

*ETH Zürich, Institute for Theoretical Physics, juerg@phys.ethz.ch.

†University of Geneva, Section of Mathematics, antti.knowles@unige.ch.

‡University of Zürich, Institute of Mathematics, benjamin.schlein@math.uzh.ch.

§University of Warwick, Mathematics Institute, V.Sohinger@warwick.ac.uk.

Given \mathbb{P} as in (1.1), natural objects to consider are the associated *time-dependent correlation functions*. More precisely, for $m \in \mathbb{N}$, times $t_1, \dots, t_m \in \mathbb{R}$, and functions $X^1, \dots, X^m \in C^\infty(\mathfrak{H})$, we consider

$$Q_{\mathbb{P}}(X^1, \dots, X^m; t_1, \dots, t_m) := \int X^1(S_{t_1}u) \cdots X^m(S_{t_m}u) d\mathbb{P}(u), \quad (1.2)$$

the m -particle time-dependent correlation function associated with H . The goal of this paper is a microscopic derivation of (1.2) from the corresponding many-body quantum objects in the case when the Hamiltonian flow is the flow of a cubic nonlinear Schrödinger equation in one spatial dimension. This is the time-dependent variant of the question previously considered in [11, 18], and subsequently in [19, 20].

We now set this up more precisely. Let us consider the spatial domain $\Lambda = \mathbb{T}^1$ or \mathbb{R} . The *one-particle space* is given by $\mathfrak{H} := L^2(\Lambda; \mathbb{C})$. The scalar product and norm on \mathfrak{H} are denoted by $\langle \cdot, \cdot \rangle_{\mathfrak{H}}$ and $\| \cdot \|_{\mathfrak{H}}$ respectively. We use the convention that $\langle \cdot, \cdot \rangle_{\mathfrak{H}}$ is linear in the second argument. We start from the *one-body Hamiltonian*

$$h := -\Delta + \kappa + v, \quad (1.3)$$

for a *chemical potential* $\kappa > 0$ and a *one-body potential* $v : \Lambda \rightarrow [0, +\infty)$. This is a positive, self-adjoint densely defined operator on \mathfrak{H} . Furthermore, we assume that h has a compact resolvent and satisfies

$$\mathrm{Tr} h^{-1} < \infty. \quad (1.4)$$

In particular, we can take $v = 0$ when $\Lambda = \mathbb{T}^1$. We write the spectral representation of h as

$$h = \sum_{k \in \mathbb{N}} \lambda_k u_k u_k^*. \quad (1.5)$$

Here $\lambda_k > 0$ are the eigenvalues and u_k are the associated normalized eigenfunctions in \mathfrak{H} of the operator h . We consider an *interaction potential* w that satisfies

$$w \in L^\infty(\Lambda), \quad w \geq 0 \text{ pointwise}. \quad (1.6)$$

The Hamilton function that we consider is

$$H(u) := \int_{\Lambda} dx (|\nabla u(x)|^2 + v(x)|u(x)|^2) + \frac{1}{2} \int_{\Lambda \times \Lambda} dx dy |u(x)|^2 w(x-y) |u(y)|^2, \quad (1.7)$$

where dx denotes the Lebesgue measure on Λ . We often abbreviate $\int_{\Lambda} dx \equiv \int dx$. The space of fields $u : \Lambda \rightarrow \mathbb{C}$ generates a Poisson algebra where the Poisson bracket is given by

$$\{u(x), \bar{u}(y)\} = i\delta(x-y), \quad \{u(x), u(y)\} = \{\bar{u}(x), \bar{u}(y)\} = 0. \quad (1.8)$$

The Hamiltonian equation of motion associated with (1.7)–(1.8) is the nonlocal *nonlinear Schrödinger equation (NLS)*

$$i\partial_t u(x) + (\Delta - \kappa)u(x) = v(x)u(x) + \int_{\Lambda} dy |u(y)|^2 w(x-y)u(x). \quad (1.9)$$

In addition to (1.9), we also consider the local NLS

$$i\partial_t u(x) + (\Delta - \kappa)u(x) = |u(x)|^2 u(x), \quad (1.10)$$

obtained from (1.9) by setting $v = 0$ and $w = \delta$. This is the Hamiltonian equation of motion associated with the Hamiltonian obtained from (1.7) by the analogous modifications.

By the arguments of [3] we know that both (1.9) and (1.10) are globally well-posed in \mathfrak{H} . Given initial data $u_0 \in \mathfrak{H}$, we denote the solution at time t by

$$u(t) =: S_t u_0. \quad (1.11)$$

1.1. The quantum problem. We use the same conventions as in [11, Section 1.4]. We work on the *bosonic Fock space*

$$\mathcal{F} \equiv \mathcal{F}(\mathfrak{H}) := \bigoplus_{p \in \mathbb{N}} \mathfrak{H}^{(p)}.$$

Here, for $p \in \mathbb{N}$, the p -particle space $\mathfrak{H}^{(p)}$ is defined as the symmetric subspace of $\mathfrak{H}^{\otimes p}$. For $f \in \mathfrak{H}$ let $b^*(f)$ and $b(f)$ denote the usual bosonic creation and annihilation operators on \mathcal{F} , defined by

$$(b^*(f)\Psi)^{(p)}(x_1, \dots, x_p) = \frac{1}{\sqrt{p}} \sum_{i=1}^p f(x_i) \Psi^{(p-1)}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_p), \quad (1.12)$$

$$(b(f)\Psi)^{(p)}(x_1, \dots, x_p) = \sqrt{p+1} \int dx \bar{f}(x) \Psi^{(p+1)}(x, x_1, \dots, x_p), \quad (1.13)$$

where we denote vectors of \mathcal{F} by $\Psi = (\Psi^{(p)})_{p \in \mathbb{N}}$. They satisfy the canonical commutation relations

$$[b(f), b^*(g)] = \langle f, g \rangle_{\mathfrak{H}}, \quad [b(f), b(g)] = [b^*(f), b^*(g)] = 0. \quad (1.14)$$

We define the rescaled creation and annihilation operators

$$\phi_\tau^*(f) := \tau^{-1/2} b^*(f), \quad \phi_\tau(f) := \tau^{-1/2} b(f). \quad (1.15)$$

We think of ϕ_τ^* and ϕ_τ as operator-valued distributions and we denote their distribution kernels as $\phi_\tau^*(x)$ and $\phi_\tau(x)$ respectively. In analogy to the classical field ϕ defined in (1.22) below, we call ϕ_τ the quantum field. For more details, we refer the reader to [11, Section 1.4].

Let $p \in \mathbb{N}$ and ξ a closed linear operator on $\mathfrak{H}^{(p)}$, given by a Schwartz integral kernel that we denote by $\xi(x_1, \dots, x_p; y_1, \dots, y_p)$; see [23, Corollary V.4.4]. We define the *lift* of ξ to \mathcal{F} by

$$\Theta_\tau(\xi) := \int dx_1 \cdots dx_p dy_1 \cdots dy_p \xi(x_1, \dots, x_p; y_1, \dots, y_p) \phi_\tau^*(x_1) \cdots \phi_\tau^*(x_p) \phi_\tau(y_1) \cdots \phi_\tau(y_p). \quad (1.16)$$

The *quantum interaction* is defined as

$$\mathcal{W}_\tau := \frac{1}{2} \Theta_\tau(W) = \frac{1}{2} \int dx dy \phi_\tau^*(x) \phi_\tau^*(y) w(x-y) \phi_\tau(x) \phi_\tau(y). \quad (1.17)$$

Here $W \equiv W^{(2)}$ is the two particle operator on $\mathfrak{H}^{(2)}$ given by multiplication by $w(x_1 - x_2)$ for w as in (1.6). The *free quantum Hamiltonian* is given by

$$H_{\tau,0} := \Theta_\tau(h) = \int dx dy \phi_\tau^*(x) h(x; y) \phi_\tau(y), \quad (1.18)$$

where $h(x; y)$ denotes the operator kernel of the operator h defined in (1.3).

The *interacting quantum Hamiltonian* is defined as

$$H_\tau := H_{\tau,0} + \mathcal{W}_\tau. \quad (1.19)$$

The *grand canonical ensemble* is defined as $P_\tau := e^{-H_\tau}$. We define the *quantum state* $\rho_\tau(\cdot)$ as

$$\rho_\tau(\mathcal{A}) := \frac{\text{Tr}(\mathcal{A}P_\tau)}{\text{Tr}(P_\tau)} \quad (1.20)$$

for \mathcal{A} a closed operator on \mathcal{F} .

Remark 1.1. We give a heuristic motivation for our choice of scaling (1.15), which ensures that both the terms in (1.19) have a nontrivial limit as $\tau \rightarrow \infty$. We consider the case $w = 0$. Then, it is shown in [11, (1.29)] that for $f, g \in \mathfrak{H}$, we have

$$\lim_{\tau \rightarrow \infty} \rho_\tau(\phi_\tau^*(f)\phi_\tau(g)) = \langle f, h^{-1}g \rangle.$$

One obtains a similar statement for arbitrary polynomials in ϕ_τ and ϕ_τ^* by using Wick's theorem. Thus, at least in the noninteracting case $w = 0$, any polynomial in the rescaled operators ϕ_τ and ϕ_τ^* has a well-defined expectation in the state $\rho_\tau(\cdot)$ as $\tau \rightarrow \infty$. As a consequence, the scaling in (1.15) ensures that the quadratic $H_{\tau,0}$ and the quartic \mathcal{W}_τ are typically of the same order in the state $\rho_\tau(\cdot)$. More physically, this scaling can be understood by noting that the expected number of particles $\tau\mathcal{N}_\tau$ (see (2.1) below for the definition of the operator \mathcal{N}_τ) in the state $\rho_\tau(\cdot)$ is of order τ .

In what follows, it is helpful to work with the *rescaled* version of the interacting quantum Hamiltonian given by τH_τ .

Definition 1.2. Let \mathbf{A} be an operator on the Fock space \mathcal{F} . We define its quantum time evolution as

$$\Psi_\tau^t \mathbf{A} := e^{it\tau H_\tau} \mathbf{A} e^{-it\tau H_\tau}.$$

1.2. The classical problem. For each $k \in \mathbb{N}$, let μ_k be a standard complex Gaussian measure, i.e. $\mu_k(dz) = \frac{1}{\pi} e^{-|z|^2} dz$, where dz is the Lebesgue measure on \mathbb{C} . We then introduce the probability space $(\mathbb{C}^{\mathbb{N}}, \mathcal{G}, \mu)$, with \mathcal{G} the product sigma-algebra and the product probability measure

$$\mu := \bigotimes_{k \in \mathbb{N}} \mu_k. \quad (1.21)$$

Elements of the corresponding probability space $\mathbb{C}^{\mathbb{N}}$ are denoted by $\omega = (\omega_k)_{k \in \mathbb{N}}$.

We denote by $\phi \equiv \phi(\omega)$ the *free classical field*

$$\phi := \sum_{k \in \mathbb{N}} \frac{\omega_k}{\sqrt{\lambda_k}} u_k. \quad (1.22)$$

Note that, by (1.4), the sum (1.22) converges in \mathfrak{H} almost surely.

For a closed operator ξ on $\mathfrak{H}^{(p)}$, in analogy to (1.16), we define the random variable

$$\Theta(\xi) := \int dx_1 \cdots dx_p dy_1 \cdots dy_p \xi(x_1, \dots, x_p; y_1, \dots, y_p) \bar{\phi}(x_1) \cdots \bar{\phi}(x_p) \phi(y_1) \cdots \phi(y_p). \quad (1.23)$$

Note that if ξ is a bounded operator then $\Theta(\xi)$ is almost surely well-defined, since $\phi \in \mathfrak{H}$ almost surely.

Given w as in (1.6), the *classical interaction* is defined as

$$\mathcal{W} := \frac{1}{2}\Theta(W) = \frac{1}{2} \int dx dy |\phi(x)|^2 w(x-y) |\phi(y)|^2. \quad (1.24)$$

Moreover, the *free classical Hamiltonian* is given by

$$H_0 := \Theta(h) = \int dx dy \bar{\phi}(x) h(x; y) \phi(y). \quad (1.25)$$

The *interacting classical Hamiltonian* is given by

$$H := H_0 + \mathcal{W}. \quad (1.26)$$

We define the *classical state* $\rho(\cdot)$ as

$$\rho(X) := \frac{\int X e^{-\mathcal{W}} d\mu}{\int e^{-\mathcal{W}} d\mu}, \quad (1.27)$$

where X is a random variable.

Definition 1.3. Let $p \in \mathbb{N}$ and ξ be a bounded operator on $\mathfrak{H}^{(p)}$. We define the random variable

$$\Psi^t \Theta(\xi) := \int dx_1 \cdots dx_p dy_1 \cdots dy_p \xi(x_1, \dots, x_p; y_1, \dots, y_p) \overline{S_t \phi}(x_1) \cdots \overline{S_t \phi}(x_p) S_t \phi(y_1) \cdots S_t \phi(y_p),$$

where S_t is the flow map from (1.11). Note that $\Psi^t \Theta(\xi)$ is well defined since $\phi \in \mathfrak{H}$ almost surely and since S_t preserves the norm on \mathfrak{H} .

1.3. Statement of the main results. We denote by $\mathcal{L}(\mathcal{H})$ the space of bounded operators on a Hilbert space \mathcal{H} . We prove the following result for the flow of (1.9).

Theorem 1.4 (Convergence of time-dependent correlation functions for the nonlocal nonlinearity). *Given $m \in \mathbb{N}$, $p_1, \dots, p_m \in \mathbb{N}$, $\xi^1 \in \mathcal{L}(\mathfrak{H}^{(p_1)}), \dots, \xi^m \in \mathcal{L}(\mathfrak{H}^{(p_m)})$ and $t_1, \dots, t_m \in \mathbb{R}$, we have*

$$\lim_{\tau \rightarrow \infty} \rho_\tau \left(\Psi_\tau^{t_1} \Theta_\tau(\xi^1) \cdots \Psi_\tau^{t_m} \Theta_\tau(\xi^m) \right) = \rho \left(\Psi^{t_1} \Theta(\xi^1) \cdots \Psi^{t_m} \Theta(\xi^m) \right).$$

Remark 1.5. For all $p \in \mathbb{N}$, $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$, and $t \in \mathbb{R}$ we have by (1.20), Definition 1.2, and the cyclicity of the trace that

$$\rho_\tau \left(\Psi_\tau^t \Theta_\tau(\xi) \right) = \rho_\tau \left(\Theta_\tau(\xi) \right) \quad (1.28)$$

for all τ . In particular, substituting (1.28) into Theorem 1.4 with $m = 1$, it follows that

$$\rho \left(\Psi^t \Theta(\xi) \right) = \rho \left(\Theta(\xi) \right). \quad (1.29)$$

Hence, using (1.28)–(1.29), we recover the invariance of the Gibbs measure for (1.9), proved in [4].

Choosing physical space to be a circle, $\Lambda = \mathbb{T}^1$, and the external potential to vanish, $v = 0$, we prove an analogue of Theorem 1.4 for the dynamics corresponding to a local nonlinearity (see (1.10)) by using an approximation argument. Let w be a continuous compactly supported nonnegative function satisfying $\int dx w(x) = 1$. For $\varepsilon > 0$ we define the two-body potential

$$w^\varepsilon(x) := \frac{1}{\varepsilon} w \left(\frac{[x]}{\varepsilon} \right). \quad (1.30)$$

Here, and in the sequel, $[x]$ denotes the unique element of the set $(x + \mathbb{Z}) \cap [-1/2, 1/2)$.

Theorem 1.6 (Convergence of time-dependent correlation functions for a local non-linearity). *Suppose that $\Lambda = \mathbb{T}^1$, $v = 0$, and w^ε is defined as in (1.30). There exists a sequence (ε_τ) of positive numbers satisfying $\lim_{\tau \rightarrow \infty} \varepsilon_\tau = 0$, such that, for arbitrary $m \in \mathbb{N}$, $p_1, \dots, p_m \in \mathbb{N}$, $\xi^1 \in \mathcal{L}(\mathfrak{H}^{(p_1)}), \dots, \xi^m \in \mathcal{L}(\mathfrak{H}^{(p_m)})$, and $t_1 \in \mathbb{R}, \dots, t_m \in \mathbb{R}$, we have*

$$\lim_{\tau \rightarrow \infty} \rho_\tau^{\varepsilon_\tau} \left(\Psi_\tau^{t_1, \varepsilon_\tau} \Theta_\tau(\xi^1) \dots \Psi_\tau^{t_m, \varepsilon_\tau} \Theta_\tau(\xi^m) \right) = \rho \left(\Psi^{t_1} \Theta(\xi^1) \dots \Psi^{t_m} \Theta(\xi^m) \right).$$

Here, the quantum state $\rho_\tau^\varepsilon(\cdot)$ is defined in (1.20) and the quantum-mechanical time evolution $\Psi_\tau^{t, \varepsilon}$ is introduced in Definition 1.2, where the two-body potential is w^ε . Moreover, the classical state $\rho(\cdot)$ is defined in (1.27) and the classical time evolution Ψ^t is introduced in Definition 1.3, where the two-body potential is $w = \delta$. (Hence the classical time evolution is governed by the local nonlinear Schrödinger equation (1.10)).

Remark 1.7. As in Remark 1.5, Theorem 1.6 allows us to establish the invariance of the Gibbs measure for (1.10) first proved in [4].

Remark 1.8. For interacting Bose gases on a finite lattice, results similar to Theorems 1.4 and 1.6 have been obtained in [16, Section 3.4].

Conventions. We write $\mathbb{N} = \{0, 1, 2, \dots\}$. We denote by C a positive constant that can depend on the fixed quantities of the problem (for example the interaction potential w). This constant can change from line to line. If it depends on a family of parameters a_1, a_2, \dots , we write $C = C(a_1, a_2, \dots)$. Given a separable Hilbert space \mathcal{H} and $q \in [1, \infty]$, we denote by $\mathfrak{S}^q(\mathcal{H})$ the q -Schatten class. This is the set of all $\mathcal{T} \in \mathcal{L}(\mathcal{H})$ such that the norm given by

$$\|\mathcal{T}\|_{\mathfrak{S}^q(\mathcal{H})} := \begin{cases} (\text{Tr} |\mathcal{T}|^q)^{1/q} & \text{if } q < \infty \\ \sup \text{spec} |\mathcal{T}| & \text{if } q = \infty \end{cases}$$

is finite. Here we recall that $|\mathcal{T}| := \sqrt{\mathcal{T}^* \mathcal{T}}$. In particular, we note that by definition $\mathfrak{S}^\infty(\mathcal{H}) = \mathcal{L}(\mathcal{H})$, the space of bounded operators on \mathcal{H} . We abbreviate the operator norm $\|\cdot\|_{\mathfrak{S}^\infty}$ by $\|\cdot\|$. Any quantity bearing a subscript τ is a quantum object and any quantity not bearing this subscript is a classical object.

Acknowledgements. We would like to thank the referee for their helpful comments on our earlier version of the manuscript.

2. Strategy of the proof

We first outline the strategy of proof of Theorem 1.4, concerning the nonlocal problem. Let us recall several definitions. The *rescaled number of particles* is defined as

$$\mathcal{N}_\tau := \int dx \phi_\tau^*(x) \phi_\tau(x). \quad (2.1)$$

Moreover, the *mass* is defined as

$$\mathcal{N} := \int dx |\phi(x)|^2. \quad (2.2)$$

Theorem 1.4 can be deduced from the following two propositions.

Proposition 2.1 (Convergence in the small particle number regime). *Let $F \in C_c^\infty(\mathbb{R})$ with $F \geq 0$ be given. Given $m \in \mathbb{N}$, $p_1, \dots, p_m \in \mathbb{N}$, $\xi^1 \in \mathcal{L}(\mathfrak{H}^{(p_1)}), \dots, \xi^m \in \mathcal{L}(\mathfrak{H}^{(p_m)})$, and $t_1, \dots, t_m \in \mathbb{R}$, we have*

$$\lim_{\tau \rightarrow \infty} \rho_\tau \left(\Psi_\tau^{t_1} \Theta_\tau(\xi^1) \cdots \Psi_\tau^{t_m} \Theta_\tau(\xi^m) F(\mathcal{N}_\tau) \right) = \rho \left(\Psi^{t_1} \Theta(\xi^1) \cdots \Psi^{t_m} \Theta(\xi^m) F(\mathcal{N}) \right).$$

Proposition 2.2 (Bounds in the large particle number regime). *Let $G \in C^\infty(\mathbb{R})$ be such that $0 \leq G \leq 1$ and $G = 0$ on $[0, \mathcal{K}]$ for some $\mathcal{K} > 0$. Furthermore, let $m \in \mathbb{N}$, $p_1, \dots, p_m \in \mathbb{N}$, $\xi^1 \in \mathcal{L}(\mathfrak{H}^{(p_1)}), \dots, \xi^m \in \mathcal{L}(\mathfrak{H}^{(p_m)})$, and $t_1, \dots, t_m \in \mathbb{R}$ be given. The following estimates hold.*

$$(i) \left| \rho_\tau \left(\Psi_\tau^{t_1} \Theta_\tau(\xi^1) \cdots \Psi_\tau^{t_m} \Theta_\tau(\xi^m) G(\mathcal{N}_\tau) \right) \right| \leq \frac{C}{\mathcal{K}}.$$

$$(ii) \left| \rho \left(\Psi^{t_1} \Theta(\xi^1) \cdots \Psi^{t_m} \Theta(\xi^m) G(\mathcal{N}) \right) \right| \leq \frac{C}{\mathcal{K}}.$$

Here $C = C(\|\xi^1\|, \dots, \|\xi^m\|, p_1 + \dots + p_m) > 0$ is a constant that does not depend on \mathcal{K} .

Proof of Theorem 1.4. For fixed $\mathcal{K} > 0$, we choose $F \equiv F_\mathcal{K}$ in Proposition 2.1 such that $0 \leq F \leq 1$ and $F = 1$ on $[0, \mathcal{K}]$ and we let $G \equiv G_\mathcal{K} := 1 - F_\mathcal{K}$ in Proposition 2.2. We then deduce Theorem 1.4 by letting $\mathcal{K} \rightarrow \infty$. \square

We prove Proposition 2.1 in Section 3 and Proposition 2.2 in Section 4 below.

Theorem 1.6, concerning the local problem, is proved in Section 5 by using Theorem 1.4 and a limiting argument. At this step, it is important to prove an L^2 -convergence result of solutions of the NLS with interaction potential given by (1.30) to solutions of (1.10). The precise statement is given in Proposition 5.1 below. Note that, in order to prove this statement, it is not enough to use energy methods, but we have to directly use the dispersion in the problem. To this end, we use $X^{s,b}$ spaces, which are recalled in Definition 5.2 below.

3. The small particle number regime: proof of Proposition 2.1.

In this section we consider the small particle number regime and prove Proposition 2.1.

In what follows, it is useful to note that, given $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$, for $\Theta_\tau(\xi)$ defined as in (1.16), we have

$$\Theta_\tau(\xi)|_{\mathfrak{H}^{(n)}} = \begin{cases} \frac{p!}{\tau^p} \binom{n}{p} P_+(\xi \otimes \mathbf{1}^{(n-p)}) P_+ & \text{if } n \geq p \\ 0 & \text{otherwise.} \end{cases} \quad (3.1)$$

(For more details see [16, (3.88)].) Here $\mathbf{1}^{(q)}$ denotes the identity map on $\mathfrak{H}^{(q)}$ and P_+ denotes the orthogonal projection onto the subspace of symmetric tensors. In particular (c.f. [16, Section 3.4.1]), we deduce the following estimate.

Lemma 3.1. *Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$ be given. For all $n \in \mathbb{N}$ we have*

$$\left\| \Theta_\tau(\xi)|_{\mathfrak{H}^{(n)}} \right\| \leq \left(\frac{n}{\tau} \right)^p \|\xi\|.$$

Moreover, by applying the Cauchy-Schwarz inequality, we obtain the following result in the classical setting (c.f. also [Section 3.4.2] [16]).

Lemma 3.2. *Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$ be given. Then we have*

$$|\Theta(\xi)| \leq \|\phi\|_{\mathfrak{H}}^{2p} \|\xi\|.$$

3.1. An auxiliary convergence result. In the proof of Proposition 2.1, we use the following auxiliary convergence result.

For $p \in \mathbb{N}$ define the unit ball $\mathfrak{B}_p := \{\eta \in \mathfrak{S}^2(\mathfrak{H}^{(p)}) : \|\eta\|_{\mathfrak{S}^2(\mathfrak{H}^{(p)})} \leq 1\}$.

Proposition 3.3. *Let $f \in C_c^\infty(\mathbb{R})$ be given.*

(i) *We have*

$$\lim_{\tau \rightarrow \infty} \rho_\tau(\Theta_\tau(\xi)f(\mathcal{N}_\tau)) = \rho(\Theta(\xi)f(\mathcal{N})), \quad (3.2)$$

uniformly in $\xi \in \mathfrak{B}_p \cup \{\mathbf{1}^{(p)}\}$.

(ii) *Moreover, if $f \geq 0$ then (3.2) holds for all $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$.*

We note that, if f were equal to 1 (which is not allowed in the assumptions), then the result of Proposition 3.3 would follow immediately from [18, Theorem 5.3] or equivalently [11, Theorem 1.8]. However, Proposition 3.3 does not immediately follow from the arguments in [11] since the presence of f breaks the Gaussian structure which allows us to apply the Wick theorem. In the proof we expand f by means of complex analytic methods in such a way that we can apply the analysis of [11] in the result.

Before we proceed with the proof of Proposition 3.3, we introduce some notation and collect several auxiliary results.

For \mathcal{N} as in (2.2) and $\nu > 0$ we define the measure

$$d\tilde{\mu}^\nu := e^{-\nu\mathcal{N}} d\mu.$$

Note that $d\tilde{\mu}^\nu$ is still Gaussian, but it is not normalized. Indeed, recalling (1.21), we write

$$d\mu(\omega) = \bigotimes_{k \in \mathbb{N}} \frac{1}{\pi} e^{-|\omega_k|^2} d\omega_k.$$

We have

$$\mathcal{N} = \int dx |\phi(x)|^2 = \sum_{k \in \mathbb{N}} \frac{|\omega_k|^2}{\lambda_k},$$

and so we find

$$d\tilde{\mu}^\nu(\omega) = \bigotimes_{k \in \mathbb{N}} \frac{1}{\pi} e^{-\nu|\omega_k|^2/\lambda_k} e^{-|\omega_k|^2} d\omega_k. \quad (3.3)$$

Define the normalized Gaussian measure

$$d\mu^\nu := \frac{d\tilde{\mu}^\nu}{\int d\tilde{\mu}^\nu}.$$

The measure μ^ν satisfies a Wick theorem, where any moment of variables that are linear functions of ϕ or $\bar{\phi}$ is given as a sum over pairings and each pair is computed using the (Hermitian) covariance of μ^ν given by

$$h^\nu := h + \nu = \sum_{k \in \mathbb{N}} (\lambda_k + \nu) u_k u_k^*. \quad (3.4)$$

In terms of ϕ we have

$$\int d\mu^\nu \langle \phi, g \rangle \langle f, \phi \rangle = \langle f, (h^\nu)^{-1} g \rangle.$$

For $\operatorname{Re} z \geq 0$ and for X a random variable and for \mathcal{W} as in (1.24), we define the deformed classical state

$$\tilde{\rho}_z^\nu(X) := \int X e^{-z\mathcal{W}} d\mu^\nu. \quad (3.5)$$

In the quantum setting, for $\operatorname{Re} z \geq 0$ and for \mathcal{A} a closed operator on \mathcal{F} , we define the deformed quantum state

$$\tilde{\rho}_{\tau,z}^\nu(\mathcal{A}) := \frac{\operatorname{Tr}(\mathcal{A} e^{-H_{\tau,0} - z\mathcal{W}_\tau - \nu\mathcal{N}_\tau})}{\operatorname{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}. \quad (3.6)$$

The free state $\tilde{\rho}_{\tau,0}^\nu(\cdot)$ satisfies a quantum Wick theorem (c.f. [11, Appendix B]), with the quantum Green function

$$G_\tau^\nu := \frac{1}{\tau(e^{h^\nu/\tau} - 1)}.$$

In the proof of Proposition 3.3 we have to analyse

$$\frac{\operatorname{Tr}(\mathcal{A} e^{-H_{\tau,0} - z\mathcal{W}_\tau - \nu\mathcal{N}_\tau})}{\operatorname{Tr}(e^{-H_{\tau,0}})} = \tilde{\rho}_{\tau,z}^\nu(\mathcal{A}) \frac{\operatorname{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}{\operatorname{Tr}(e^{-H_{\tau,0}})}.$$

With the above notation we have the following result.

Lemma 3.4. *For $\nu > 0$ we have*

$$\lim_{\tau \rightarrow \infty} \frac{\operatorname{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}{\operatorname{Tr}(e^{-H_{\tau,0}})} = \int d\tilde{\mu}^\nu.$$

Proof of Lemma 3.4. A direct calculation using (3.3) shows that

$$\int d\tilde{\mu}^\nu = \prod_{k \in \mathbb{N}} \frac{\lambda_k}{\lambda_k + \nu}. \quad (3.7)$$

As in the proof of [11, Appendix B, Proof of Lemma B.1], we use the *occupation state basis*. This is an orthonormal basis $(\psi_{\mathbf{m}})_{\mathbf{m} \in \mathbb{N}^{\mathbb{N}}}$, of \mathcal{F} given by $\psi_{\mathbf{m}} := \prod_{k \in \mathbb{N}} \frac{(b_k^*)^{m_k}}{\sqrt{m_k!}} \Omega$, where $\Omega = (1, 0, 0, \dots) \in \mathcal{F}$ is the vacuum state and $\mathbf{m} = (m_k)_{k \in \mathbb{N}}$ ranges over $\mathbb{N}^{\mathbb{N}}$. Using the observation that $b_l \Omega = 0$ for all $l \in \mathbb{N}$ and the canonical commutation relations (1.14) we obtain

$$\begin{cases} b_l \psi_{\mathbf{m}} = \sqrt{m_l} \psi_{\mathbf{m}'} & \text{for } m'_k = m_k - \delta_{kl} \\ b_l^* \psi_{\mathbf{m}} = \sqrt{m_l + 1} \psi_{\mathbf{m}'} & \text{for } m'_k = m_k + \delta_{kl}. \end{cases} \quad (3.8)$$

Applying (1.5), (1.18), (2.1), and (3.8) we deduce that

$$\frac{\operatorname{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}{\operatorname{Tr}(e^{-H_{\tau,0}})} = \frac{\sum_{\mathbf{m}} e^{-\sum_k \frac{\lambda_k + \nu}{\tau} m_k}}{\sum_{\mathbf{m}} e^{-\sum_k \frac{\lambda_k}{\tau} m_k}} = \frac{\sum_{\mathbf{m}} \prod_k e^{-\frac{\lambda_k + \nu}{\tau} m_k}}{\sum_{\mathbf{m}} \prod_k e^{-\frac{\lambda_k}{\tau} m_k}} = \prod_{k \in \mathbb{N}} \frac{1 - e^{-\frac{\lambda_k}{\tau}}}{1 - e^{-\frac{\lambda_k + \nu}{\tau}}}. \quad (3.9)$$

We note that, for fixed $k \in \mathbb{N}$, we have

$$\frac{1 - e^{-\frac{\lambda_k}{\tau}}}{1 - e^{-\frac{\lambda_k + \nu}{\tau}}} = 1 + \frac{e^{-\frac{\lambda_k}{\tau}} (e^{-\frac{\nu}{\tau}} - 1)}{1 - e^{-\frac{\lambda_k + \nu}{\tau}}} = 1 + \mathcal{O}\left(\frac{\nu}{\lambda_k}\right). \quad (3.10)$$

Indeed, we note that by the mean value theorem

$$e^{-\frac{\nu}{\tau}} - 1 = \mathcal{O}\left(\frac{\nu}{\tau}\right). \quad (3.11)$$

Also, we observe that

$$\left|1 - e^{-\frac{\lambda_k + \nu}{\tau}}\right| \geq \left|1 - e^{-\frac{\lambda_k}{\tau}}\right|. \quad (3.12)$$

If $\lambda_k < \tau$ we have

$$\left|1 - e^{-\frac{\lambda_k}{\tau}}\right| \geq \frac{C\lambda_k}{\tau}, \quad e^{-\frac{\lambda_k}{\tau}} \leq 1. \quad (3.13)$$

Furthermore, if $\lambda_k \geq \tau$ we have

$$\left|1 - e^{-\frac{\lambda_k}{\tau}}\right| \geq C, \quad e^{-\frac{\lambda_k}{\tau}} \leq \frac{C\tau}{\lambda_k}. \quad (3.14)$$

The estimate (3.10) follows from (3.11)–(3.14).

Note that the individual factors of (3.9) converge to the corresponding factors of (3.7) as $\tau \rightarrow \infty$. We hence reduce the claim to showing that

$$\lim_{\tau \rightarrow \infty} \prod_{\substack{k \in \mathbb{N}: \\ \lambda_k \gg \nu}} \frac{1 - e^{-\frac{\lambda_k}{\tau}}}{1 - e^{-\frac{\lambda_k + \nu}{\tau}}} = \prod_{\substack{k \in \mathbb{N}: \\ \lambda_k \gg \nu}} \frac{\lambda_k}{\lambda_k + \nu}. \quad (3.15)$$

Here we use the notation $A \gg B$ if there exists a large constant $C > 0$ such that $A \geq CB$. (The size of C is specified from context). The convergence (3.15) follows from the dominated convergence theorem after taking logarithms on both sides, using (3.10), the inequality $|\log(1+z)| \leq C|z|$ for $|z| \leq 1/2$ and the assumption that $\text{Tr } h^{-1} < \infty$. Taking logarithms is justified by (3.10) and the assumption $\lambda_k \gg \nu$. \square

Before proceeding to the proof of Proposition 3.3 we recall the *Helffer-Sjöstrand formula*.

Proposition 3.5 (Helffer-Sjöstrand formula). *Let $g \in C^2(\mathbb{R})$ be given. We define the almost-analytic extension of g of degree 1 as a function on \mathbb{C} given by*

$$\tilde{g}(u + iv) := g(u) + ivg'(u).$$

Let $\chi \in C_c^\infty(\mathbb{C}; [0, 1])$ be given. Then, for any $\lambda \in \mathbb{R}$ with $\chi(\lambda) = 1$ we have

$$g(\lambda) = \frac{1}{\pi} \int_{\mathbb{C}} d\zeta \frac{\partial_{\bar{\zeta}}(\tilde{g}(\zeta) \chi(\zeta))}{\lambda - \zeta}$$

Above we use the notation $\zeta = u + iv$ for $u = \text{Re } \zeta$ and $v = \text{Im } \zeta$. In particular, the antiholomorphic derivative is given by $\partial_{\bar{\zeta}} = \frac{1}{2}(\partial_u + i\partial_v)$. Furthermore, $d\zeta$ denotes Lebesgue measure on \mathbb{C} .

For a proof of Proposition 3.5, we refer the reader for instance to [1, Proposition C.1] with $n = 1$. We also recall a result that gives us conditions under which we can deduce convergence in the trace class from convergence in the Hilbert-Schmidt norm.

Lemma 3.6. *Let $p \in \mathbb{N}$ be given. Suppose that for all $\tau > 0$, $\gamma_\tau \in \mathfrak{S}^1(\mathfrak{H}^{(p)})$ is positive and $\gamma \in \mathfrak{S}^1(\mathfrak{H}^{(p)})$ is positive (as an operator). Moreover, suppose that we have*

$$(i) \quad \lim_{\tau \rightarrow \infty} \|\gamma_\tau - \gamma\|_{\mathfrak{S}^2(\mathfrak{H}^{(p)})} = 0, \quad (ii) \quad \lim_{\tau \rightarrow \infty} \text{Tr } \gamma_\tau = \text{Tr } \gamma.$$

Then we have $\lim_{\tau \rightarrow \infty} \|\gamma_\tau - \gamma\|_{\mathfrak{S}^1(\mathfrak{H}^{(p)})} = 0$.

Lemma 3.6 was proved in [11, Lemma 4.10], based on arguments given in [25, Section 2] (in particular, on the proof of [25, Lemma 2.20]). We now have all of the necessary ingredients to prove Proposition 3.3.

Proof of Proposition 3.3. We first prove (i). Let us consider

$$\xi \in \mathcal{C}_p := \mathcal{B}_p \cup \{\mathbf{1}^{(p)}\}.$$

For $\zeta \in \mathbb{C} \setminus [0, \infty)$, we define the functions $\alpha_\tau^\xi \equiv \alpha_\tau^\xi(\zeta)$ and $\alpha^\xi \equiv \alpha^\xi(\zeta)$ by

$$\alpha_\tau^\xi(\zeta) := \rho_\sharp \left(\Theta_\sharp(\xi) \frac{1}{\mathcal{N}_\sharp - \zeta} \right). \quad (3.16)$$

In the above formula and throughout the proof of the proposition, we use the convention that, given $Y = \mathcal{N}, \alpha, \rho, \dots$, the quantity Y_\sharp formally denotes either Y_τ or Y . In this convention, we write ϕ^* for $\bar{\phi}$. This simplifies some of the notation in the sequel.

For $\text{Re } \zeta < 0$ we have

$$\frac{1}{\mathcal{N}_\sharp - \zeta} = \int_0^\infty d\nu e^{-(\mathcal{N}_\sharp - \zeta)\nu} = \int_0^\infty d\nu e^{\zeta\nu} e^{-\nu\mathcal{N}_\sharp}. \quad (3.17)$$

In particular, from (3.16)–(3.17), it follows that for $\text{Re } \zeta < 0$ we have

$$\alpha_\tau^\xi(\zeta) = \int_0^\infty d\nu e^{\zeta\nu} \rho_\sharp(\Theta_\sharp(\xi) e^{-\nu\mathcal{N}_\sharp}). \quad (3.18)$$

By Lemma 3.1 we know that $\pm\Theta_\tau(\xi) \leq \|\xi\| \mathcal{N}_\tau^p \leq \mathcal{N}_\tau^p$ acting on sectors of Fock space (c.f. [16, (3.91)]). Hence, it follows that

$$|\rho_\tau(\Theta_\tau(\xi) e^{-\nu\mathcal{N}_\tau})| \leq \rho_\tau(\mathcal{N}_\tau^p e^{-\nu\mathcal{N}_\tau}) \leq \rho_\tau(\mathcal{N}_\tau^p) \leq C(p), \quad (3.19)$$

uniformly in $\xi \in \mathcal{C}_p$ and $\nu > 0$. Furthermore, by using Lemma 3.2, we deduce the classical analogue of (3.19),

$$|\rho(\Theta(\xi) e^{-\nu\mathcal{N}})| \leq \rho(\mathcal{N}^p e^{-\nu\mathcal{N}}) \leq \rho(\mathcal{N}^p) \leq C(p), \quad (3.20)$$

uniformly in $\nu > 0$. The estimates (3.19)–(3.20) and the assumption $\text{Re } \zeta < 0$ allow us to use Fubini's theorem in order to exchange the integration in ν and expectation $\rho_\sharp(\cdot)$ in (3.16)–(3.17) and deduce (3.18). For fixed $\nu > 0$ we have

$$\begin{aligned} \rho_\tau(\Theta_\tau(\xi) e^{-\nu\mathcal{N}_\tau}) &= \frac{\text{Tr}(\Theta_\tau(\xi) e^{-H_{\tau,0} - \mathcal{W}_\tau - \nu\mathcal{N}_\tau})}{\text{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})} \frac{\text{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}{\text{Tr}(e^{-H_{\tau,0}})} \frac{\text{Tr}(e^{-H_{\tau,0}})}{\text{Tr}(e^{-H_\tau})} \\ &= \tilde{\rho}_{\tau,1}^\nu(\Theta_\tau(\xi)) \frac{\text{Tr}(e^{-H_{\tau,0} - \nu\mathcal{N}_\tau})}{\text{Tr}(e^{-H_{\tau,0}})} \frac{1}{\tilde{\rho}_{\tau,1}^0(1)}, \end{aligned} \quad (3.21)$$

where $\tilde{\rho}_{\tau,z}^0$ is defined by setting $\nu = 0$ in (3.6). Moreover, we have

$$\rho(\Theta(\xi)e^{-\nu\mathcal{N}}) = \left(\int d\mu^\nu \Theta(\xi) e^{-W} \right) \left(\int d\tilde{\mu}^\nu \right) \left(\frac{1}{\int d\mu e^{-W}} \right) = \tilde{\rho}_1^\nu(\Theta(\xi)) \left(\int d\tilde{\mu}^\nu \right) \frac{1}{\tilde{\rho}_1^0(1)}, \quad (3.22)$$

where $\tilde{\rho}_z^0$ is defined by setting $\nu = 0$ in (3.5).

We now consider each of the factors in (3.21)–(3.22). By [11, Theorem 1.8] with the Hamiltonian h^ν given by (3.4), the first factor in (3.21) converges to the first factor in (3.22) as $\tau \rightarrow \infty$ uniformly in $\xi \in \mathcal{C}_p$. Moreover, the convergence of the second factors follows from Lemma 3.4. Finally, we have convergence of the third factors by [11, Theorem 1] with the Hamiltonian h . Note that both applications of [11, Theorem 1.8] are justified by the assumption (1.4). In particular, we obtain that

$$\lim_{\tau \rightarrow \infty} \rho_\tau(\Theta_\tau(\xi)e^{-\nu\mathcal{N}_\tau}) = \rho(\Theta(\xi)e^{-\nu\mathcal{N}}), \quad (3.23)$$

uniformly in $\xi \in \mathcal{C}_p$. From (3.18), (3.19), (3.23) and the dominated convergence theorem, it follows that for $\text{Re } \zeta < 0$ we have

$$\lim_{\tau \rightarrow \infty} \alpha_\tau^\xi(\zeta) = \alpha^\xi(\zeta), \quad (3.24)$$

uniformly in $\xi \in \mathcal{C}_p$.

By (3.16), Hölder's inequality, and arguing as in (3.19) and (3.20) (with $\nu = 0$), we have

$$|\alpha_\tau^\xi(\zeta)| \leq \left\| \frac{1}{\mathcal{N}_\tau - \zeta} \right\| \rho_\tau(\mathcal{N}_\tau^p) \leq \frac{C(p)}{|\text{Im } \zeta|}. \quad (3.25)$$

We now show that $\alpha_\tau^\xi, \alpha^\xi$ are analytic in $\mathbb{C} \setminus [0, \infty)$.

In the sequel, we use the notation

$$\mathfrak{H}^{(\leq R)} := \bigoplus_{p \leq R} \mathfrak{H}^{(p)}, \quad \mathfrak{H}^{(\geq R)} := \bigoplus_{p \geq R} \mathfrak{H}^{(p)} \quad (3.26)$$

for $R > 0$. Let us denote the corresponding orthogonal projections by

$$P^{(\leq R)} : \mathcal{F} \rightarrow \mathfrak{H}^{(\leq R)}, \quad P^{(\geq R)} : \mathcal{F} \rightarrow \mathfrak{H}^{(\geq R)}. \quad (3.27)$$

In order to prove the analyticity of α_τ^ξ in $\mathbb{C} \setminus [0, \infty)$ we argue similarly as in the proof of [11, Lemma 2.34]. Namely, given $n \in \mathbb{N}$ we define for $\zeta \in \mathbb{C} \setminus [0, \infty)$

$$\alpha_{\tau,n}^\xi(\zeta) := \rho_\tau \left(P^{(\leq n)} \Theta_\tau(\xi) \frac{1}{\mathcal{N}_\tau - \zeta} \right).$$

Here $P^{(\leq n)}$ is defined as in (3.27). Note that $\alpha_{\tau,n}^\xi$ is analytic in $\mathbb{C} \setminus [0, \infty)$ since \mathcal{N}_τ is constant on each m -particle sector of \mathcal{F} . As in (3.25) we note that

$$|\alpha_{\tau,n}^\xi(\zeta)| \leq C(p) \left\| \frac{1}{\mathcal{N}_\tau - \zeta} \right\| \leq \frac{C(p)}{\max\{-\text{Re } \zeta, |\text{Im } \zeta|\}}.$$

Finally, for $\zeta \in \mathbb{C} \setminus [0, \infty)$ we know that $\lim_{n \rightarrow \infty} \alpha_{\tau,n}^\xi(\zeta) = \alpha_\tau^\xi(\zeta)$ by construction. The analyticity of $\alpha_\tau^\xi(\zeta)$ in $\mathbb{C} \setminus [0, \infty)$ now follows. The analyticity of α^ξ in $\mathbb{C} \setminus [0, \infty)$ is verified by using Lemma 3.2 and differentiating under the integral sign in the representation

$$\alpha^\xi(\zeta) = \frac{\int d\mu \Theta(\xi) \frac{1}{\mathcal{N} - \zeta} e^{-W}}{\int d\mu e^{-W}}.$$

In what follows, we define the function $\beta_\tau^\xi : \mathbb{C} \setminus [0, \infty) \rightarrow \mathbb{C}$ by

$$\beta_\tau^\xi := \alpha_\tau^\xi - \alpha^\xi. \quad (3.28)$$

From the analyticity of α_τ^ξ on $\mathbb{C} \setminus [0, \infty)$, (3.24) and (3.25), we note that the β_τ^ξ satisfy the following properties.

- (1) β_τ^ξ is analytic on $\mathbb{C} \setminus [0, \infty)$.
- (2) $\lim_{\tau \rightarrow \infty} \sup_{\xi \in \mathcal{C}_p} |\beta_\tau^\xi(\zeta)| = 0$ for all $\operatorname{Re} \zeta < 0$.
- (3) $\sup_{\xi \in \mathcal{C}_p} |\beta_\tau^\xi(\zeta)| \leq \frac{C(p)}{|\operatorname{Im} \zeta|}$ for all $\zeta \in \mathbb{C} \setminus [0, \infty)$.

We now show that

$$\lim_{\tau \rightarrow \infty} \sup_{\xi \in \mathcal{C}_p} |\beta_\tau^\xi(\zeta)| = 0 \quad \text{for all } \zeta \in \mathbb{C} \setminus [0, \infty). \quad (3.29)$$

Namely, we generalise condition (2) above to all $\zeta \in \mathbb{C} \setminus [0, \infty)$.

Given $\varepsilon > 0$ we define

$$\mathcal{D}_\varepsilon := \{\zeta : \operatorname{Im} \zeta > \varepsilon\}$$

and

$$\mathcal{T}_\varepsilon := \{\zeta_0 \in \mathcal{D}_\varepsilon : \lim_{\tau \rightarrow \infty} \sup_{\xi \in \mathcal{C}_p} |\partial_\zeta^m \beta_\tau^\xi(\zeta_0)| \rightarrow 0 \text{ for all } m \in \mathbb{N}\}.$$

In other words, \mathcal{T}_ε consists of all points in \mathcal{D}_ε at which all ζ -derivatives of β_τ^ξ converge to zero as $\tau \rightarrow \infty$, uniformly in $\xi \in \mathcal{C}_p$. Note that, by using conditions (1)-(3) above, Cauchy's integral formula and the dominated convergence theorem we have $\mathcal{D}_\varepsilon \cap \{\zeta : \operatorname{Re} \zeta < 0\} \subset \mathcal{T}_\varepsilon$ hence $\mathcal{T}_\varepsilon \neq \emptyset$.

In order to prove (3.29) on \mathcal{D}_ε , it suffices to show that $\mathcal{T}_\varepsilon = \mathcal{D}_\varepsilon$. By connectedness of \mathcal{D}_ε and since $\mathcal{T}_\varepsilon \neq \emptyset$, the latter claim follows if we prove that \mathcal{T}_ε is both open and closed in \mathcal{D}_ε . Let us first prove that \mathcal{T}_ε is open in \mathcal{D}_ε . Given $\zeta_0 \in \mathcal{T}_\varepsilon$, we note that $B_{\zeta_0}(\varepsilon/2) \subset \mathcal{D}_{\varepsilon/2}$. Hence by property (3), it follows that $|\beta_\tau^\xi| \leq C(\varepsilon)$ on $B_{\zeta_0}(\varepsilon/2)$. By analyticity and Cauchy's integral formula it follows that the series expansion of β_τ^ξ at ζ_0 converges on $B_{\zeta_0}(\varepsilon/2)$. Therefore, by differentiating term by term and using the dominated convergence theorem and the assumption $\zeta_0 \in \mathcal{T}_\varepsilon$, it follows that $B_{\zeta_0}(\varepsilon/2) \subset \mathcal{T}_\varepsilon$. Hence, \mathcal{T}_ε is open in \mathcal{D}_ε .

We now show that \mathcal{T}_ε is closed in \mathcal{D}_ε . Suppose that (ζ_n) is a sequence in \mathcal{T}_ε such that $\zeta_n \rightarrow \zeta$ for some $\zeta \in \mathcal{D}_\varepsilon$. We now show that $\zeta \in \mathcal{T}_\varepsilon$. In order to do this, we note that for n large enough we have $\zeta \in B_{\zeta_n}(\varepsilon/2)$. The argument used to show the openness of \mathcal{T}_ε in \mathcal{D}_ε gives us that $B_{\zeta_n}(\varepsilon/2) \subset \mathcal{T}_\varepsilon$. In particular, $\zeta \in \mathcal{T}_\varepsilon$. Hence \mathcal{T}_ε is closed in \mathcal{D}_ε . Therefore $\mathcal{T}_\varepsilon = \mathcal{D}_\varepsilon$. The convergence (3.29) is shown on $\tilde{\mathcal{D}}_\varepsilon := \{\zeta : \operatorname{Im} \zeta < -\varepsilon\}$ by symmetry. The claim (3.29) on all of $\mathbb{C} \setminus [0, \infty)$ now follows by letting $\varepsilon \rightarrow 0$ and recalling that (3.29) holds for $\zeta < 0$ by condition (2) above.

By Proposition 3.5 we obtain

$$f(\mathcal{N}_\sharp) = \frac{1}{\pi} \int_{\mathbb{C}} d\zeta \frac{\partial_{\bar{\zeta}}[(f(u) + ivf'(u))\chi(v)]}{\mathcal{N}_\sharp - \zeta}, \quad (3.30)$$

where $\chi \in C_c^\infty(\mathbb{R})$ is a function such that $\chi = 1$ on $[-1, 1]$. More precisely, we use the assumption that $f \in C_c^\infty(\mathbb{R})$ in order to deduce that we can take χ to be a compactly supported function in the v variable. Furthermore, χ can be taken to be equal to 1 on $[-1, 1]$ since $\operatorname{spec}(\mathcal{N}_\tau) \subset \mathbb{R}$ (c.f. [1, (C.1)]) and since \mathcal{N} takes values in \mathbb{R} .

Let us define

$$\psi(\zeta) := \frac{1}{\pi} \partial_{\bar{\zeta}} [(f(u) + iv f'(u)) \chi(v)] = \frac{1}{2\pi} [iv f''(u) \chi(v) + i(f(u) + iv f'(u)) \chi'(v)],$$

so that by (3.30) we have

$$f(\mathcal{N}_{\#}) = \int_{\mathbb{C}} d\zeta \frac{\psi(\zeta)}{\mathcal{N}_{\#} - \zeta}. \quad (3.31)$$

Since $f, \chi \in C_c^\infty(\mathbb{R})$, it follows that

$$\psi \in C_c^\infty(\mathbb{C}). \quad (3.32)$$

By our choice of ψ we know that

$$|\psi(\zeta)| \leq C|v| = C|\operatorname{Im} \zeta|. \quad (3.33)$$

Substituting this into (3.31) we deduce that

$$\rho_{\#}(\Theta_{\#}(\xi) f(\mathcal{N}_{\#})) = \int_{\mathbb{C}} d\zeta \psi(\zeta) \rho_{\#} \left(\Theta_{\#}(\xi) \frac{1}{\mathcal{N}_{\#} - \zeta} \right) = \int_{\mathbb{C}} d\zeta \psi(\zeta) \alpha_{\#}^{\xi}(\zeta). \quad (3.34)$$

We note that, by (3.25), (3.32) and (3.33) we have for almost all $\zeta \in \mathbb{C}$

$$|\psi(\zeta) \alpha_{\#}^{\xi}(\zeta)| \leq F(\zeta) \quad (3.35)$$

for some function $F \in L^1(\mathbb{C})$. Therefore, the interchanging of the integration in ζ and expectation $\rho_{\#}(\cdot)$ in (3.34) is justified by Fubini's theorem. Furthermore, recalling (3.28) and using (3.29), we note that

$$\lim_{\tau \rightarrow \infty} \sup_{\xi \in \mathcal{C}_p} |\alpha_{\#}^{\xi}(\zeta) - \alpha^{\xi}(\zeta)| = 0 \text{ for all } \zeta \in \mathbb{C} \setminus [0, \infty). \quad (3.36)$$

The claim (i) now follows from (3.34)–(3.36) and the dominated convergence theorem.

We now prove (ii). Let us define $\gamma_{\#,p}^f$ by duality according to

$$\operatorname{Tr} (\gamma_{\#,p}^f \eta) = \rho_{\#}(\Theta_{\#}(\eta) f(\mathcal{N}_{\#})),$$

for $\eta \in \mathcal{L}(\mathfrak{H}^{(p)})$. In particular

$$\gamma_{\#,p}^f(x_1, \dots, x_p; y_1, \dots, y_p) = \rho_{\#}(\phi_{\#}^*(y_1) \cdots \phi_{\#}^*(y_p) \phi_{\#}(x_1) \cdots \phi_{\#}(x_p) f(\mathcal{N}_{\#})). \quad (3.37)$$

By duality, part (i) implies that $\lim_{\tau \rightarrow \infty} \|\gamma_{\#,p}^f - \gamma_p^f\|_{\mathfrak{S}^2(\mathfrak{H}^{(p)})} = 0$ and $\lim_{\tau \rightarrow \infty} \operatorname{Tr} \gamma_{\#,p}^f = \operatorname{Tr} \gamma_p^f$. The claim follows from Lemma 3.6 if we prove that the $\gamma_{\#,p}^f$ are positive operators. Namely, if this is the case, Lemma 3.6 implies that we have $\lim_{\tau \rightarrow \infty} \|\gamma_{\#,p}^f - \gamma_p^f\|_{\mathfrak{S}^1(\mathfrak{H}^{(p)})} = 0$ and claim (ii) then follows by duality.

We now prove the positivity of $\gamma_{\#,p}^f$. Given $\eta \in \mathfrak{H}^{(p)}$, a direct calculation using (3.37) shows that we have

$$\langle \eta, \gamma_{\#,p}^f \eta \rangle_{\mathfrak{H}^{(p)}} = \rho_{\#}(\Theta_{\#}(\eta \otimes \bar{\eta}) f(\mathcal{N}_{\#})).$$

This quantity is nonnegative in the quantum setting since $\Theta_{\#}(\eta \otimes \bar{\eta}), f(\mathcal{N}_{\#}), e^{-H_{\tau,0} - \mathcal{W}_{\tau}}$ are positive operators on \mathcal{F} . In order to see the positivity of $\Theta_{\#}(\eta \otimes \bar{\eta})$, we apply (3.1). Moreover, in the classical setting, the quantities

$$\Theta(\eta \otimes \bar{\eta}) = \left| \int dx_1 \cdots dx_p \bar{\phi}(x_1) \cdots \bar{\phi}(x_p) \eta(x_1, \dots, x_p) \right|^2, \quad f(\mathcal{N}), \quad e^{-W}$$

are nonnegative. Therefore the $\gamma_{\#,p}^f$ are indeed positive operators. Note that this is the only step where we use the nonnegativity of f . \square

3.2. Schwinger-Dyson expansion in the quantum problem. Arguing similarly as in [16, Section 4.2], we apply a Schwinger-Dyson expansion to $\Psi_\tau^t \Theta_\tau(\xi)$. Here we recall the time-evolution operator Ψ_τ^t from Definition 1.2. We note that a related approach was also applied in [10, 12].

Before we proceed with the expansion, we first introduce the operation \bullet_r as well as the free quantum time evolution of operators on \mathcal{F} , analogously to Definition 1.2.

Definition 3.7. Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$, $\eta \in \mathcal{L}(\mathfrak{H}^{(q)})$ and $r \leq \min\{p, q\}$ be given.

(i) We define

$$\xi \bullet_r \eta := P_+(\xi \otimes \mathbf{1}^{(q-r)})(\mathbf{1}^{(p-r)} \otimes \eta)P_+ \in \mathcal{L}(\mathfrak{H}^{(p+q-r)}),$$

where we recall that P_+ denotes the orthogonal projection from $\mathfrak{H}^{\otimes r}$ to $\mathfrak{H}^{(r)}$.

(ii) With \bullet_r given by (i), we define

$$[\xi, \eta]_r := \xi \bullet_r \eta - \eta \bullet_r \xi \in \mathcal{L}(\mathfrak{H}^{(p+q-r)}).$$

The following lemma can be found in [16, Section 3.4.1]. We omit the proof.

Lemma 3.8. Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$, $\eta \in \mathcal{L}(\mathfrak{H}^{(q)})$ and $r \leq \min\{p, q\}$ be given. The following identities hold.

$$(i) \quad \Theta_\tau(\xi)\Theta_\tau(\eta) = \sum_{r=0}^{\min\{p,q\}} \binom{p}{r} \binom{q}{r} \frac{r!}{\tau^r} \Theta_\tau(\xi \bullet_r \eta).$$

$$(ii) \quad [\Theta_\tau(\xi), \Theta_\tau(\eta)] = \sum_{r=1}^{\min\{p,q\}} \binom{p}{r} \binom{q}{r} \frac{r!}{\tau^r} \Theta_\tau([\xi, \eta]_r).$$

Definition 3.9. Let \mathbf{A} be an operator on \mathcal{F} . We define its free quantum time evolution by

$$\Psi_{\tau,0}^t \mathbf{A} := e^{it\tau H_{\tau,0}} \mathbf{A} e^{-it\tau H_{\tau,0}}.$$

Note that, using first-quantized notation, we have

$$\tau H_{\tau,0} \Big|_{\mathfrak{H}^{(n)}} = \sum_{i=1}^n h_i. \tag{3.38}$$

Here h_i denotes the operator h acting in the x_i variable. By (3.38), we note the operator $\Psi_{\tau,0}^t$ does not depend on τ . We keep the subscript τ in order to emphasize that this is a quantum time evolution. Moreover, it is useful to apply a time evolution to p -particle operators.

Definition 3.10. Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$. For $t \in \mathbb{R}$ we define

$$\xi_t := e^{it \sum_{j=1}^p h_j} \xi e^{-it \sum_{j=1}^p h_j}.$$

In particular, from (1.16), Definition 3.9, (3.38) and Definition 3.10, it follows that for $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$ we have

$$\Psi_{\tau,0}^t \Theta_\tau(\xi) = \Theta_\tau(\xi_t). \tag{3.39}$$

The following result holds.

Lemma 3.11. *Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$. Given $\mathcal{K} > 0$, $\varepsilon > 0$ and $t \in \mathbb{R}$, there exists $L = L(\mathcal{K}, \varepsilon, t, \|\xi\|, p) \in \mathbb{N}$, a finite sequence $(e^l)_{l=0}^L$ with $e^l = e^l(\xi, t) \in \mathcal{L}(\mathfrak{H}^{(l)})$ and $\tau_0 = \tau_0(\mathcal{K}, \varepsilon, t, \|\xi\|) > 0$ such that*

$$\left\| \left(\Psi_\tau^t \Theta_\tau(\xi) - \sum_{l=0}^L \Theta_\tau(e^l) \right) \Big|_{\mathfrak{H}^{(\leq \mathcal{K}\tau)}} \right\| \leq \varepsilon, \quad (3.40)$$

for all $\tau \geq \tau_0$. Note that $\mathfrak{H}^{(\leq \mathcal{K}\tau)}$ is defined as in (3.26) above.

Proof. Let us first observe that

$$\Psi_\tau^t \Theta_\tau(\xi) = \Theta_\tau(\xi_t) + (ip) \int_0^t ds \Psi_\tau^s \Theta_\tau([W, \xi_{t-s}]_1) + \frac{i\binom{p}{2}}{\tau} \int_0^t ds \Psi_\tau^s \Theta_\tau([W, \xi_{t-s}]_2). \quad (3.41)$$

Indeed, we write

$$\Psi_\tau^t \Theta_\tau(\xi) = \Psi_\tau^s \Psi_{\tau,0}^{-s} \Psi_{\tau,0}^t \Theta_\tau(\xi) \Big|_{s=t} = \Psi_{\tau,0}^t \Theta_\tau(\xi) + \int_0^t ds \frac{d}{ds} \left(\Psi_\tau^s \Psi_{\tau,0}^{-s} \Psi_{\tau,0}^t \Theta_\tau(\xi) \right), \quad (3.42)$$

which by (3.39) and Definitions 1.2 and 3.9 equals

$$\Theta_\tau(\xi_t) + \int_0^t ds \frac{d}{ds} \left(e^{is\tau H_\tau} e^{-is\tau H_{\tau,0}} \Theta_\tau(\xi_t) e^{is\tau H_{\tau,0}} e^{-is\tau H_\tau} \right). \quad (3.43)$$

By differentiating in s and using (1.19), (1.18) and (3.39), it follows that the integrand in the second term of (3.43) equals

$$\frac{i\tau}{2} \Psi_\tau^s \Psi_{\tau,0}^{-s} [\Theta_\tau(W_s), \Theta_\tau(\xi_t)] = \frac{i\tau}{2} \Psi_\tau^s [\Theta_\tau(W), \Theta_\tau(\xi_{t-s})],$$

which by Lemma 3.8 (ii) equals

$$(ip) \Theta_\tau([W, \xi_{t-s}]_1) + \frac{i\binom{p}{2}}{\tau} \Theta_\tau([W, \xi_{t-s}]_2). \quad (3.44)$$

Substituting (3.44) into (3.43), we deduce (3.41).

Iteratively applying (3.41) we deduce that, for all $M \in \mathbb{N}$ we have

$$\Psi_\tau^t \Theta_\tau(\xi) = A_{\tau,M}^t(\xi) + E_{\tau,M}^t(\xi) + B_{\tau,M}^t(\xi),$$

where

$$\begin{aligned} A_{\tau,M}^t(\xi) &:= \Theta_\tau(\xi_t) + \sum_{j=1}^{M-1} i^j p(p+1) \cdots (p+j-1) \left\{ \int_0^t ds_1 \int_0^{s_1} ds_2 \cdots \int_0^{s_{j-1}} ds_j \right. \\ &\quad \left. \Theta_\tau \left([W_{s_j}, [W_{s_{j-1}}, \dots, [W_{s_1}, \xi_t]_1 \cdots]_1]_1 \right) \right\}, \\ E_{\tau,M}^t(\xi) &:= i^M p(p+1) \cdots (p+M-1) \left\{ \int_0^t ds_1 \int_0^{s_1} ds_2 \cdots \int_0^{s_{M-1}} ds_M \right. \\ &\quad \left. \Psi_\tau^{s_M} \Theta_\tau \left([W, [W_{s_{M-1}-s_M}, \dots, [W_{s_1-s_M}, \xi_{t-s_M}]_1 \cdots]_1]_1 \right) \right\}, \\ B_{\tau,M}^t(\xi) &:= \frac{1}{\tau} \sum_{j=1}^M i^j p(p+1) \cdots (p+j-2) \binom{p+j-1}{2} \left\{ \int_0^t ds_1 \int_0^{s_1} ds_2 \cdots \int_0^{s_{j-1}} ds_j \right. \\ &\quad \left. \Psi_\tau^{s_j} \Theta_\tau \left([W, [W_{s_{j-1}-s_j}, \dots, [W_{s_1-s_j}, \xi_{t-s_j}]_1 \cdots]_1]_2 \right) \right\}. \quad (3.45) \end{aligned}$$

Moreover, we define $A_{\tau,\infty}^t(\xi)$ and $B_{\tau,\infty}^t(\xi)$ by formally setting $M = \infty$ in (3.45). We now show that, on $\mathfrak{H}^{(\leq \mathcal{K}\tau)}$ we have

$$A_{\tau,M}^t(\xi) \rightarrow A_{\tau,\infty}^t(\xi), \quad E_{\tau,M}^t(\xi) \rightarrow 0, \quad B_{\tau,M}^t(\xi) \rightarrow B_{\tau,\infty}^t(\xi) \quad (3.46)$$

as $M \rightarrow \infty$ in norm whenever $|t| < T_0(\mathcal{K})$, where $T_0(\mathcal{K})$ is chosen sufficiently small depending on \mathcal{K} , but *independent of p* . In particular, it follows that on $\mathfrak{H}^{(\leq \mathcal{K}\tau)}$, the formally-defined quantities $A_{\tau,\infty}^t(\xi)$ and $B_{\tau,\infty}^t(\xi)$ are well defined and that $E_{\tau,\infty}^t(\xi)$ vanishes.

In order to prove (3.46) we note that, if $n \leq \mathcal{K}\tau$, the j -th term of the formal sum $A_{\tau,\infty}^t(\xi)$ acting on $\mathfrak{H}^{(n)}$ is estimated in norm by

$$\frac{|t|^j}{j!} (p+j)^j 2^j \left(\frac{n}{\tau}\right)^{p+j} \|w\|_{L^\infty}^j \|\xi\|. \quad (3.47)$$

Here we used Lemma 3.1 as well as $\|\xi_t\| = \|\xi\|$, $\|W_s\| = \|W\| = \|w\|_{L^\infty}$. The latter two equalities follow immediately from Definition 3.10. The expression in (3.47) is

$$\leq e^p \mathcal{K}^p \left(2e \mathcal{K} \|w\|_{L^\infty} |t|\right)^j \|\xi\|. \quad (3.48)$$

Using (3.48), we can deduce the first convergence result in (3.46) for $|t| < T_0(\mathcal{K})$. By noting that Ψ_τ^s preserves the operator norm, we deduce the second and third convergence results in (3.46) by an analogous argument. We omit the details. In particular, on $\mathfrak{H}^{(\leq \mathcal{K}\tau)}$ we can write for $|t| < T_0(\mathcal{K})$

$$\Psi_\tau^t \Theta_\tau(\xi) = A_{\tau,\infty}^t(\xi) + B_{\tau,\infty}^t(\xi), \quad (3.49)$$

where the infinite sum converges in norm. Recalling (3.45), it also follows from this proof that

$$\|B_{\tau,\infty}^t(\xi)|_{\mathfrak{H}^{(\leq \mathcal{K}\tau)}}\| \leq \frac{Ce^p \mathcal{K}^p \|\xi\|}{\tau}. \quad (3.50)$$

By (3.45)–(3.46), (3.49)–(3.50), we deduce that (3.40) holds for $|t| < T_0(\mathcal{K})$. Note that the e^l are obtained from the partial sums of $A_{\tau,\infty}^t(\xi)$ (as in (3.45)). By construction we have that $e^l \in \mathcal{L}(\mathfrak{H}^{(l)})$.

We obtain (3.40) for general t by iterating this procedure in increments of size $T_0(\mathcal{K})$. This is possible to do by using norm conservation, i.e. we use that for all operators \mathbf{A} on \mathcal{F} we have

$$\|\Psi_\tau^t \mathbf{A}|_{\mathfrak{H}^{(\leq \mathcal{K}\tau)}}\| = \|\mathbf{A}|_{\mathfrak{H}^{(\leq \mathcal{K}\tau)}}\|. \quad (3.51)$$

Furthermore, we use the observation that the radius of convergence $T_0(\mathcal{K})$ does not depend on p . The latter fact is required since after each iteration of the procedure we generate q -particle operators, where q grows with t . A detailed description of an analogous iteration procedure applied in a slightly different context can be found in [16, Lemma 3.6]. \square

3.3. Schwinger-Dyson expansion in the classical problem. The following lemma can be found in [16, Section 3.4.2]. We omit the proof.

Lemma 3.12. *Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$, $\eta \in \mathcal{L}(\mathfrak{H}^{(q)})$ be given. We then have*

$$\{\Theta(\xi), \Theta(\eta)\} = ipq \Theta([\xi, \eta]_1).$$

Definition 3.13. Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$. We define $\Psi_0^t \Theta(\xi)$ to be the random variable

$$\int dx_1 \cdots dx_p dy_1 \cdots dy_p \xi(x_1, \dots, x_p; y_1, \dots, y_p) \overline{S_{t,0}\phi}(x_1) \cdots \overline{S_{t,0}\phi}(x_p) S_{t,0}\phi(y_1) \cdots S_{t,0}\phi(y_p),$$

where $S_{t,0} := e^{-ith}$ denotes the free Schrödinger evolution on \mathfrak{H} corresponding to the Hamiltonian (1.3).

In particular, from Definitions 1.3 and 3.13 we have

$$\partial_t \Psi^t \Theta(\xi) = \Psi^t \{H, \Theta(\xi)\}, \quad \partial_t \Psi_0^t \Theta(\xi) = \Psi_0^t \{H_0, \Theta(\xi)\}, \quad (3.52)$$

where we recall (1.26)–(1.25).

From (1.23), Definition 3.10 and Definition 3.13, it follows that for $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$ we have

$$\Psi_0^t \Theta(\xi) = \Theta(\xi_t). \quad (3.53)$$

We now prove the classical analogue of Lemma 3.11.

Lemma 3.14. *Let $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$. Given $\mathcal{K} > 0$, $\varepsilon > 0$ and $t \in \mathbb{R}$, for $L = L(\mathcal{K}, \varepsilon, t, \|\xi\|, p) \in \mathbb{N}$ chosen possibly larger than in Lemma 3.11 and for the same choice of $e^l = e^l(\xi, t) \in \mathcal{L}(\mathfrak{H}^{(l)})$ as in Lemma 3.11 we have*

$$\left| \left(\Psi^t \Theta(\xi) - \sum_{l=0}^L \Theta(e^l) \right) \mathbf{1}_{\{\mathcal{N} \leq \mathcal{K}\}} \right| \leq \varepsilon.$$

Proof. We first note that we have the classical analogue of (3.41)

$$\Psi^t \Theta(\xi) = \Theta(\xi_t) + (ip) \int_0^t ds \Psi^s \Theta([W, \xi_{t-s}]_1). \quad (3.54)$$

Namely, arguing as in (3.42) and using (3.53), it follows that

$$\Psi^t \Theta(\xi) = \Theta(\xi_t) + \int_0^t ds \frac{d}{ds} \left(\Psi^s \Psi_0^{-s} \Psi_0^t \Theta(\xi) \right). \quad (3.55)$$

Differentiating and using (3.52), it follows that the integrand in (3.55) equals

$$\Psi^s \{H, \Psi_0^{-s+t} \Theta(\xi)\} - \Psi^s \Psi_0^{-s} \{H_0, \Psi_0^t \Theta(\xi)\} = \Psi^s \{H, \Theta(\xi_{-s+t})\} - \Psi^s \Psi_0^{-s} \{H_0, \Theta(\xi_t)\}.$$

In the last equality we also used (3.53). By Lemma 3.12 and (1.26), we can rewrite this as

$$\Psi^s \left(ip \Theta([h + W, \xi_{-s+t}]_1) \right) - \Psi^s \left(\Psi_0^{-s} ip \Theta([h, \xi_t]_1) \right). \quad (3.56)$$

We note that $\Psi_0^{-s} \Theta([h, \xi_t]_1) = \Theta([h, \xi_{-t+s}]_1)$ and hence the expression in (3.56) equals

$$ip \Psi^s \Theta([W, \xi_{t-s}]_1).$$

Substituting this into (3.55) we obtain (3.54).

We now iterate (3.54) analogously as in the proof of Lemma 3.11. The convergence for $|t| < T_0(\mathcal{K})$ is shown by arguing as in the proof of (3.46). The only difference is that instead of applying Lemma 3.1, we now apply Lemma 3.2. (In fact, the quantity $T_0(\mathcal{K})$ can be chosen to be the same as the corresponding quantity in Lemma 3.11, which was obtained from (3.48)). Furthermore, in the extension to all times, instead of applying (3.51), we use that S_t preserves the norm on \mathfrak{H} . Finally, we note that the e^l that we obtain from iterating (3.54) are the same as those obtained by iterating (3.41) in the proof of Lemma 3.11. \square

3.4. Proof of Proposition 2.1. We now combine the results of Proposition 3.3, Lemma 3.11 and Lemma 3.14 in order to prove Proposition 2.1.

Proof of Proposition 2.1. By assumption, there exists $\mathcal{K} > 0$ such that $F = 0$ on (\mathcal{K}, ∞) . Let us note that, for all $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$ and for all $t \in \mathbb{R}$, the following inequalities hold in the quantum setting.

$$\left\| \Psi_\tau^t \Theta_\tau(\xi) \Big|_{\mathfrak{H}^{(\leq \mathcal{K}\tau)}} \right\| \leq \mathcal{K}^p \|\xi\|. \quad (3.57)$$

$$\frac{\|F(\mathcal{N}_\tau) e^{-H_\tau}\|_{\mathfrak{S}^1(\mathcal{F})}}{\text{Tr}(e^{-H_\tau})} \leq C, \quad (3.58)$$

for some constant $C > 0$ independent of τ . The inequality (3.57) follows from Definition 1.2, the observation that Ψ_τ^t preserves operator norm and from Lemma 3.1. The inequality (3.58) follows from $F(\mathcal{N}_\tau) e^{-H_\tau} \geq 0$ and $\frac{\text{Tr}(F(\mathcal{N}_\tau) e^{-H_\tau})}{\text{Tr}(e^{-H_\tau})} \leq C$. Furthermore, in the classical setting, the following inequalities hold.

$$|\Psi^t \Theta(\xi)| \leq \mathcal{K}^p \|\xi\|, \quad \text{whenever} \quad \|\phi\|_{\mathfrak{H}}^2 \leq \mathcal{K}. \quad (3.59)$$

$$\rho(F(\mathcal{N})) \leq C. \quad (3.60)$$

The inequality (3.59) follows from Definition 1.3, the Cauchy-Schwarz inequality and since S_t preserves the norm on \mathfrak{H} . The inequality (3.60) is immediate since $F \in C_c^\infty(\mathbb{R})$.

We now apply Hölder's inequality, (3.57)-(3.60) and Lemmas 3.11 and 3.14 with \mathcal{K} chosen as above to deduce that the claim follows if we prove that for all $q_1, \dots, q_m \in \mathbb{N}$, $\eta^1 \in \mathcal{L}(\mathfrak{H}^{(q_1)}), \dots, \eta^m \in \mathcal{L}(\mathfrak{H}^{(q_m)})$ we have

$$\lim_{\tau \rightarrow \infty} \rho_\tau(\Theta_\tau(\eta^1) \cdots \Theta_\tau(\eta^m) F(\mathcal{N}_\tau)) = \rho(\Theta(\eta^1) \cdots \Theta(\eta^m) F(\mathcal{N})). \quad (3.61)$$

Note that in the iterative application of (3.57) we use that the operator $\Psi_\tau^t \Theta_\tau(\xi)$ leaves the sectors $\mathfrak{H}^{(n)}$ of the Fock space invariant, thus allowing us to apply the estimate on $\mathfrak{H}^{(\leq \mathcal{K}\tau)}$. We can rewrite the right-hand side of (3.61) as

$$\rho(\Theta(\eta) F(\mathcal{N})), \quad (3.62)$$

where $q := q_1 + \dots + q_m$ and

$$\eta := \eta^1 \bullet_0 \cdots \bullet_0 \eta^m \in \mathcal{L}(\mathfrak{H}^{(q)}). \quad (3.63)$$

Here we recall Definition 3.7 (i).

By iteratively applying Lemma 3.8 (i) and using Hölder's inequality together with (3.57)-(3.58), it follows that the left-hand side of (3.61) equals

$$\rho_\tau(\Theta_\tau(\eta) F(\mathcal{N}_\tau)) + \mathcal{O}\left(\frac{1}{\tau}\right), \quad (3.64)$$

where η is given by (3.63) above. The convergence (3.61) now follows from (3.62)-(3.64), the assumptions on F and Proposition 3.3 (ii). \square

4. The large particle number regime: proof of Proposition 2.2.

In this section we consider the regime where $\mathcal{N}_\tau, \mathcal{N}$ are assumed to be large. The main result that we prove is Proposition 2.2.

Proof of Proposition 2.2. We first prove (i). Let us note that \mathcal{N}_τ commutes with e^{-H_τ} and with $\Psi_\tau^{t_j}(\xi^j)$ for all $j = 1, \dots, m$. Therefore, the expression that we want to estimate in (i) can be rewritten as

$$\left| \rho_\tau \left((1 + \mathcal{N}_\tau)^{-p_1} \Psi_\tau^{t_1} \Theta_\tau(\xi^1) \cdots (1 + \mathcal{N}_\tau)^{-p_m} \Psi_\tau^{t_m} \Theta_\tau(\xi^m) (1 + \mathcal{N}_\tau)^p G(\mathcal{N}_\tau) \right) \right|,$$

where we define $p := p_1 + \dots + p_m$. Using Hölder's inequality and $(1 + \mathcal{N}_\tau)^p G(\mathcal{N}_\tau) \geq 0$, this is

$$\leq \left(\prod_{j=1}^m \left\| (1 + \mathcal{N}_\tau)^{-p_j} \Psi_\tau^{t_j} \Theta_\tau(\xi^j) \right\| \right) \rho_\tau \left((1 + \mathcal{N}_\tau)^p G(\mathcal{N}_\tau) \right). \quad (4.1)$$

The j -th factor of the first expression in (4.1) equals

$$\left\| \Psi_\tau^{t_j} (1 + \mathcal{N}_\tau)^{-p_j} \Theta_\tau(\xi^j) \right\| = \left\| (1 + \mathcal{N}_\tau)^{-p_j} \Theta_\tau(\xi^j) \right\| \leq \|\xi^j\|. \quad (4.2)$$

Here we used that \mathcal{N}_τ commutes with $e^{it_\tau H_\tau}$, that $\Psi_\tau^{t_j}$ preserves operator norm and Lemma 3.1.

By construction of G we note that the second expression in (4.1) is

$$\leq \rho_\tau \left((1 + \mathcal{N}_\tau)^p \mathbf{1}(\mathcal{N}_\tau \geq \mathcal{K}) \right),$$

which by Markov's inequality is

$$\leq \frac{\rho_\tau \left((1 + \mathcal{N}_\tau)^{p+1} \right)}{\mathcal{K}} \leq \frac{C(p)}{\mathcal{K}}. \quad (4.3)$$

The above application of Markov's inequality is justified since \mathcal{N}_τ commutes with e^{-H_τ} . Claim (i) now follows from (4.1)–(4.3).

We now prove (ii) by similar arguments. Namely, we rewrite the expression that we want to estimate in (ii) as

$$\left| \rho \left((1 + \mathcal{N})^{-p_1} \Psi^{t_1} \Theta(\xi^1) \cdots (1 + \mathcal{N})^{-p_m} \Psi^{t_m} \Theta(\xi^m) (1 + \mathcal{N})^p G(\mathcal{N}) \right) \right|,$$

which is

$$\leq \left(\prod_{j=1}^m \left| (1 + \mathcal{N})^{-p_j} \Psi^{t_j} \Theta(\xi^j) \right| \right) \rho \left((1 + \mathcal{N})^p G(\mathcal{N}) \right). \quad (4.4)$$

Using the observation that S_{t_j} preserves the norm on \mathfrak{H} as well as Lemma 3.2, it follows that the j -th factor of the first term in (4.4) is bounded by $\|\xi^j\|$. We again use the properties of G and Markov's inequality to deduce that the second term in (4.4) is

$$\leq \rho \left((1 + \mathcal{N})^p \mathbf{1}(\mathcal{N} \geq \mathcal{K}) \right) \leq \frac{\rho \left((1 + \mathcal{N})^{p+1} \right)}{\mathcal{K}} \leq \frac{C(p)}{\mathcal{K}}.$$

Claim (ii) now follows as in the quantum setting. \square

Remark 4.1. Following the proofs of Proposition 2.1 and 2.2, it is immediate that the convergence in Theorem 1.4 is uniform on the set of parameters $w \in L^\infty(\Lambda)$, $t_1 \in \mathbb{R}, \dots, t_m \in \mathbb{R}$, $p_1, \dots, p_m \in \mathbb{N}$, $m \in \mathbb{N}$, satisfying

$$\max(\|w\|_{L^\infty}, |t_1|, \dots, |t_m|, p_1, \dots, p_m, \|\xi^1\|, \dots, \|\xi^m\|, m) \leq M,$$

for any fixed $M > 0$.

5. The local problem.

In this section we fix

$$\Lambda = \mathbb{T}^1 \quad \text{and} \quad v = 0.$$

Throughout this section and Appendix A, given $s \in \mathbb{R}$, we write $H^s(\Lambda)$ for the L^2 -based inhomogeneous Sobolev space of order s on Λ .

We extend the previous analysis to the setting of the local problem (1.10). In particular, we give the proof of Theorem 1.6. Before proceeding with the proof of Theorem 1.6 we prove the following stability result.

Proposition 5.1. *Let $s \geq \frac{3}{8}$ be given. Let $\phi_0 \in H^s(\Lambda)$. We consider the Cauchy problem on Λ given by*

$$\begin{cases} i\partial_t u + (\Delta - \kappa)u = |u|^2 u \\ u|_{t=0} = \phi_0. \end{cases} \quad (5.1)$$

In addition, given $\varepsilon > 0$, and recalling the definition of w^ε from (1.30) we consider

$$\begin{cases} i\partial_t u^\varepsilon + (\Delta - \kappa)u^\varepsilon = (w^\varepsilon * |u^\varepsilon|^2)u^\varepsilon \\ u^\varepsilon|_{t=0} = \phi_0. \end{cases} \quad (5.2)$$

Let u, u^ε be solutions of (5.1) and (5.2) respectively. Then, for all $T > 0$ we have

$$\lim_{\varepsilon \rightarrow 0} \|u^\varepsilon - u\|_{L^\infty_{[-T, T]} \mathfrak{H}} = 0. \quad (5.3)$$

In order to prove Proposition 5.1 we need to recall several tools from harmonic analysis. In particular, it is helpful to use periodic Strichartz estimates formulated in $X^{\sigma, b}$ spaces. In the context of dispersive PDEs, these spaces were first used in [3].

Definition 5.2. Given $f : \Lambda_x \times \mathbb{R}_t \rightarrow \mathbb{C}$ and $\sigma, b \in \mathbb{R}$ we define

$$\|f\|_{X^{\sigma, b}} = \left\| (1 + |2\pi k|)^\sigma (1 + |\eta + 2\pi k^2|)^b \tilde{f} \right\|_{L^2_{\eta^2 k^2}},$$

where

$$\tilde{f}(k, \eta) := \int_{-\infty}^{\infty} dt \int_{\Lambda} dx f(x, t) e^{-2\pi i k x - 2\pi i \eta t}$$

denotes the *spacetime Fourier transform*.

Note that, in particular, we have

$$\|f\|_{X^{\sigma,b}} \sim \|e^{-it\Delta} f\|_{H_t^b H_x^\sigma}.$$

Here we use the convention¹ that for $h : \mathbb{R}_t \rightarrow \mathbb{C}$

$$\|h\|_{H_t^b} := \left(\int d\eta (1 + |\eta|)^{2b} |\hat{h}(\eta)|^2 \right)^{1/2}.$$

We now collect several known facts about $X^{\sigma,b}$ spaces. For a more detailed discussion we refer the reader to [27][Section 2.6] and the references therein. For the remainder of this section we fix

$$b := \frac{1}{2} + \nu, \tag{5.4}$$

for $\nu > 0$ small.

Lemma 5.3. *Let $\sigma \in \mathbb{R}$ and b as in (5.4) be given. The following properties hold.*

(i) $\|f\|_{L_t^\infty H_x^\sigma} \leq C(b) \|f\|_{X^{\sigma,b}}.$

(ii) *Suppose that $\psi \in C_c^\infty(\mathbb{R})$. Then, for all $\delta \in (0, 1)$ and $\Phi \in H^\sigma$ we have*

$$\|\psi(t/\delta) e^{it\Delta} \Phi\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \|\Phi\|_{H^\sigma}.$$

(iii) *Let ψ, δ be as in (ii). Then, for all $f \in X^{\sigma,b}$ we have*

$$\|\psi(t/\delta) f\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b}}.$$

(iv) *With the same assumptions as in (iii) we have*

$$\left\| \psi(t/\delta) \int_0^t dt' e^{i(t-t')\Delta} f(t') \right\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b-1}}.$$

(v) $\|f\|_{L_{t,x}^4} \leq C \|f\|_{X^{0,3/8}}.$

(vi) $\|f\|_{X^{0,-3/8}} \leq C \|f\|_{L_{t,x}^{4/3}}.$

(vii) *Let ψ, δ be as in (ii). Then, for all $f \in X^{0,b}$ we have*

$$\|\psi(t/\delta) f\|_{L_{t,x}^4} \leq C(b, \psi) \delta^{\theta_0} \|f\|_{X^{0,b}},$$

for some $\theta_0 \equiv \theta_0(b) > 0$.

For completeness we present a self-contained proof of Lemma 5.3 in Appendix A.

We also recall the following characterization of homogeneous Sobolev spaces on the torus.

¹We do not introduce additional factors of 2π in the definition of $\|h\|_{H_t^b}$ for simplicity of notation in the sequel.

Lemma 5.4. For $\sigma \in (0, 1)$ we have

$$\left\| \frac{f(x) - f(y)}{[x - y]^{\sigma + \frac{1}{2}}} \right\|_{L^2_{x,y}} \sim \|f\|_{\dot{H}^\sigma}. \quad (5.5)$$

Here $\|f\|_{\dot{H}^\sigma} = \|\ |\nabla|^\sigma f \|_{L^2}$ denotes the homogeneous L^2 -based Sobolev (semi)norm of order σ .

The quantity on the left-hand side of (5.5) is the periodic analogue of the *Sobolev-Slobodeckij norm*. This is a general fact. A self-contained proof using the Plancherel theorem can be found in [2, Proposition 1.3]. We now have all the tools to prove Proposition 5.1.

Proof of Proposition 5.1. We note that, in the proof, we can formally take $\kappa = 0$ for simplicity of notation. Indeed, if we let $\tilde{u} := e^{i\kappa t} u$, then \tilde{u} solves (5.1) with $\kappa = 0$. Likewise $\tilde{u}^\varepsilon := e^{i\kappa t} u^\varepsilon$ solves (5.2) with $\kappa = 0$. Finally, we note that (5.3) is equivalent to showing that for all $T > 0$ we have

$$\lim_{\varepsilon \rightarrow 0} \|\tilde{u}^\varepsilon - \tilde{u}\|_{L^\infty_{[-T, T]} \mathfrak{H}} = 0.$$

Throughout the proof, we fix $T > 0$ and consider $|t| \leq T$. In what follows, we assume $t \geq 0$. The negative times are treated by an analogous argument. Moreover we can always assume that $s < 1$. Before we proceed, we briefly recall the arguments from [26, Section 2.6] (which, in turn, are based on the arguments from [4]) used to construct the local in time solutions to (5.1) and (5.2) in H^s . Note that in [26], the quintic NLS was considered. The arguments for the cubic NLS are analogous. In what follows, we outline the main idea and refer the interested reader to the aforementioned reference for more details.

We are looking for *global mild solutions* u to (5.1)–(5.2), i.e. we want u and u^ε to solve

$$u(\cdot, t) = e^{it\Delta} \phi_0 - i \int_0^t dt' e^{i(t-t')\Delta} |u|^2 u(t') \quad (5.6)$$

$$u^\varepsilon(\cdot, t) = e^{it\Delta} \phi_0 - i \int_0^t dt' e^{i(t-t')\Delta} (w^\varepsilon * |u^\varepsilon|^2) u^\varepsilon(t') \quad (5.7)$$

for almost every t . In what follows, we construct solutions of (5.6)–(5.7) by constructing mild solutions on a sequence of intervals of fixed length depending on the initial data. Putting these solutions together, we get u and u^ε .

Let $\chi, \psi \in C_c^\infty(\mathbb{R})$ be functions such that

$$\chi(t) = \begin{cases} 1 & \text{if } |t| \leq 1 \\ 0 & \text{if } |t| > 2 \end{cases} \quad (5.8)$$

and

$$\psi(t) := \chi(t/2). \quad (5.9)$$

Given $\delta \in (0, 1)$, we define

$$\chi_\delta(t) := \chi\left(\frac{t}{\delta}\right), \quad \psi_\delta(t) := \psi\left(\frac{t}{\delta}\right). \quad (5.10)$$

Let us fix $\delta \in (0, 1)$ small which we determine later. For $t \in [0, T]$ we consider the map

$$\begin{aligned} (Lv)(\cdot, t) &:= \chi_\delta(t) e^{it\Delta} \phi_0 - i \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} |v|^2 v(t') \\ &= \chi_\delta(t) e^{it\Delta} \phi_0 - i \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} |v_\delta|^2 v_\delta(t'), \end{aligned} \quad (5.11)$$

where we define the operation

$$v_\delta(x, t) := \psi_\delta(t) v(x, t). \quad (5.12)$$

In the last equality in (5.11), we used (5.8)–(5.10). Applying Lemma 5.3 (ii)–(vii) and arguing as in the proof of [26, (2.159)] it follows that

$$\|Lv\|_{X^{s,b}} \leq C_1 \delta^{\frac{1-2b}{2}} \|\phi_0\|_{H^s} + C \delta^{r_0} \|v\|_{X^{0,b}}^2 \|v\|_{X^{s,b}} \quad (5.13)$$

$$\|Lv\|_{X^{0,b}} \leq C_1 \delta^{\frac{1-2b}{2}} \|\phi_0\|_{\mathfrak{H}} + C \delta^{r_0} \|v\|_{X^{0,b}}^3, \quad (5.14)$$

where $C_1 > 0$ is the constant from Lemma 5.3 (ii) corresponding to the cutoff in time given by χ_δ and

$$r_0 := \frac{1-2b}{2} + 3\theta_0 > 0, \quad (5.15)$$

for $\theta_0 > 0$ given by (A.42) below ². Note that, from Lemma 5.3 (ii) we know that $C_1 = C_1(\chi)$.

For clarity, we summarize the ideas of the proof of (5.14). The proof of (5.13) follows similarly using a duality argument and by applying the fractional Leibniz rule. The latter is rigorously justified by observing that the $X^{\sigma,b}$ norms are invariant under taking absolute values in the spacetime Fourier transform. For precise details on the latter point, we refer the reader to [26, (2.147)–(2.153)].

The estimate for the linear term in (5.14) follows immediately from Lemma 5.3 (ii) with $\sigma = 0$. Note that, when we apply Lemma 5.3 (iv) with $\sigma = 0$ for the Duhamel term on the right-hand side of (5.11) we have $b-1 < \frac{3}{8}$ and hence we can use Lemma 5.3 (vi), Hölder's inequality and Lemma 5.3 (v) to deduce that we have to estimate

$$\| |v_\delta|^2 v_\delta \|_{L_{t,x}^{4/3}} \leq \|v_\delta\|_{L_{t,x}^4}^3 \leq C \|v_\delta\|_{X^{0,3/8}}^3.$$

We then deduce the estimate for the Duhamel term in (5.14) by using Lemma 5.14 (vii).

Analogously, with $r_0 > 0$ as in (5.15), we have

$$\|Lv_1 - Lv_2\|_{X^{0,b}} \leq C \delta^{r_0} (\|v_1\|_{X^{0,b}}^2 + \|v_2\|_{X^{0,b}}^2) \|v_1 - v_2\|_{X^{0,b}}. \quad (5.16)$$

In particular, it follows from (5.13)–(5.16) that L is a contraction on $(\Gamma, \|\cdot\|_{X^{0,b}})$, for

$$\Gamma := \left\{ v, \|v\|_{X^{s,b}} \leq 2C_1 \delta^{\frac{1-2b}{2}} \|\phi_0\|_{H^s}, \|v\|_{X^{0,b}} \leq 2C_1 \delta^{\frac{1-2b}{2}} \|\phi_0\|_{\mathfrak{H}} \right\}, \quad (5.17)$$

where $\delta \in (0, 1)$ is chosen to be *sufficiently small depending on* $\|\phi_0\|_{\mathfrak{H}}$. By arguing as in the proof of [26, Proposition 2.3.2] (whose proof, in turn, is based on that of [9, Theorem 1.2.5]), it follows that $(\Gamma, \|\cdot\|_{X^{0,b}})$ is a closed subspace of the Banach space $X^{0,b}$, and thus complete. Therefore, we obtain a unique fixed point of L in Γ . We refer the reader to [26, Section 2.5] for more details.

²Note that the exact value of r_0 is not relevant. The main point is that it is positive. This is ensured by taking b sufficiently close to $1/2$.

Moreover, suppose that for some other $\tilde{\delta} > 0$ the function $\tilde{v} \in X^{0,b}$ solves

$$\tilde{v} = \chi_{\tilde{\delta}}(t) e^{it\Delta} \phi_0 - i \chi_{\tilde{\delta}}(t) \int_0^t dt' e^{i(t-t')\Delta} |\tilde{v}|^2 \tilde{v}(t').$$

We then want to argue that

$$v|_{\Lambda \times [0, \hat{\delta}]} = \tilde{v}|_{\Lambda \times [0, \hat{\delta}]} \quad \text{for all } \hat{\delta} \in [0, \min\{\delta, \tilde{\delta}\}]. \quad (5.18)$$

In order to prove (5.18), we need to work in *local* $X^{\sigma,b}$ spaces. Given $\sigma \in \mathbb{R}$ and a time interval I , we define

$$\|f\|_{X_I^{\sigma,b}} := \inf \left\{ \|g\|_{X^{\sigma,b}}, g|_{\Lambda \times I} = f|_{\Lambda \times I} \right\}.$$

In particular, we have that $v, \tilde{v} \in X_{[0, \hat{\delta}]}^{0,b}$. Noting that for $t \in [0, \hat{\delta}]$ we have $\chi_{\delta}(t) = \chi_{\tilde{\delta}}(t)$, the same arguments used to show (5.16) imply that

$$\|v - \tilde{v}\|_{X_{[0, \hat{\delta}]}^{0,b}} \leq C \hat{\delta}^{r_0} \left(\|v\|_{X_{[0, \hat{\delta}]}^{0,b}} + \|\tilde{v}\|_{X_{[0, \hat{\delta}]}^{0,b}} \right)^2 \|v - \tilde{v}\|_{X_{[0, \hat{\delta}]}^{0,b}} \leq C \hat{\delta}^{r_0} \left(\|v\|_{X^{0,b}} + \|\tilde{v}\|_{X^{0,b}} \right)^2 \|v - \tilde{v}\|_{X_{[0, \hat{\delta}]}^{0,b}}.$$

Here $r_0 > 0$ is given by (5.15). We hence deduce (5.18) for sufficiently small $\hat{\delta}$. By an additional iteration argument, we deduce (5.18) for the full range $\hat{\delta} \in [0, \min\{\delta, \tilde{\delta}\}]$.

Likewise, for $\varepsilon > 0$, we consider the map

$$(L^\varepsilon v)(\cdot, t) := \chi_\delta(t) e^{it\Delta} \phi_0 - i \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} (w^\varepsilon * |v_\delta|^2) v_\delta(t'), \quad (5.19)$$

for v_δ given as in (5.12). We note that, for all $k \in \mathbb{N}$, we have

$$\widehat{w^\varepsilon}(k) = \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\varepsilon} w\left(\frac{x}{\varepsilon}\right) e^{-2\pi i k x} dx = \int_{-\frac{1}{2\varepsilon}}^{\frac{1}{2\varepsilon}} w(y) e^{-2\pi i \varepsilon k y} dy.$$

Therefore, by the assumptions on w , it follows that

$$|\widehat{w^\varepsilon}(k)| \leq C \quad (5.20)$$

for some $C > 0$ independent of k, ε . Using the same arguments as for L and applying (5.20), it follows that L^ε defined in (5.19) has a unique fixed point $v^\varepsilon \in \Gamma$. Moreover, a statement analogous to (5.18) holds.

We then define u, u^ε on $[0, \delta]$ according to

$$u|_{\Lambda \times [0, \delta]} := v|_{\Lambda \times [0, \delta]}, \quad u^\varepsilon|_{\Lambda \times [0, \delta]} := v^\varepsilon|_{\Lambda \times [0, \delta]}. \quad (5.21)$$

We now iterate this construction. In doing so, we note that the increment $\delta \in (0, 1)$ we chose above *depends only on the L^2 norm of the initial data* and hence is the same at every step of the iteration. More precisely, for all $n \in \mathbb{N}$ with $n \leq (T-1)/\delta$, we construct $v_{(n)}$, and $v_{(n)}^\varepsilon$ such that the following properties hold.

- (i) $v_{(n)}$ is a mild solution of the local NLS (1.10) with $\kappa = 0$ in the sense of (5.6) on the time interval $[n\delta, (n+1)\delta]$ and we have

$$\|v_{(n)}\|_{X^{s,b}} \leq 2C_1 \delta^{\frac{1-2b}{2}} \|v_{(n)}(n\delta)\|_{H^s}. \quad (5.22)$$

- (ii) $v_{(n)}^\varepsilon$ is a mild solution of (5.2) with $\kappa = 0$ in the sense of (5.7) on the time interval $[n\delta, (n+1)\delta]$ and we have

$$\|v_{(n)}^\varepsilon\|_{X^{s,b}} \leq 2C_1 \delta^{\frac{1-2b}{2}} \|v_{(n)}^\varepsilon(n\delta)\|_{H^s}. \quad (5.23)$$

We then generalize (5.21) by defining u, u^ε on $[n\delta, (n+1)\delta]$ according to

$$u|_{\Lambda \times [n\delta, (n+1)\delta]} := v_{(n)}|_{\Lambda \times [n\delta, (n+1)\delta]}, \quad u^\varepsilon|_{\Lambda \times [n\delta, (n+1)\delta]} := v_{(n)}^\varepsilon|_{\Lambda \times [n\delta, (n+1)\delta]}. \quad (5.24)$$

Note that, in this definition, $v_{(0)} = v$ and $v_{(0)}^\varepsilon = v^\varepsilon$.

We observe that by (5.18) and the analogous uniqueness statement for L^ε , this construction does not depend on δ (as long as δ is chosen to be small enough, depending in $\|\phi_0\|_{\mathfrak{H}}$, c.f. (5.13)–(5.17)). In particular, we can choose $\delta = \delta_0(\|\phi_0\|_{\mathfrak{H}})$. By (5.24), Lemma 5.3 (i), (5.22)–(5.23), it follows that

$$\begin{cases} \|u\|_{L_{[0,T]}^\infty H_x^s} \leq C(\|\phi_0\|_{H^s}, T) \\ \|u^\varepsilon\|_{L_{[0,T]}^\infty H_x^s} \leq C(\|\phi_0\|_{H^s}, T), \end{cases} \quad (5.25)$$

for some finite quantity $C(\|\phi_0\|_{H^s}, T) > 0$. It is important to note that $C(\|\phi_0\|_{H^s}, T)$ is independent of δ . In other words, if we choose δ smaller, then the same bounds in (5.25) hold.

Using (5.21) and Lemma 5.3 (i), it follows that, for $\delta \in (0, 1)$ chosen sufficiently small as earlier, we have

$$\|u - u^\varepsilon\|_{L_{[0,\delta]}^\infty L_x^2} = \|v - v^\varepsilon\|_{L_{[0,\delta]}^\infty L_x^2} \leq C \|v - v^\varepsilon\|_{X^{0,b}}.$$

By construction of v and v^ε , we obtain

$$\begin{aligned} \|v - v^\varepsilon\|_{X^{0,b}} &\leq \left\| \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} (|v_\delta(t')|^2 - w^\varepsilon * |v_\delta(t')|^2) v_\delta(t') \right\|_{X^{0,b}} \\ &\quad + \left\| \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} \left\{ w^\varepsilon * (|v_\delta(t')|^2 - |v_\delta^\varepsilon(t')|^2) \right\} v_\delta(t') \right\|_{X^{0,b}} \\ &\quad + \left\| \chi_\delta(t) \int_0^t dt' e^{i(t-t')\Delta} \left\{ w^\varepsilon * |v_\delta^\varepsilon(t')|^2 \right\} (v_\delta(t') - v_\delta^\varepsilon(t')) \right\|_{X^{0,b}}, \end{aligned}$$

which by Lemma 5.3 (iv) is

$$\begin{aligned} &\leq C \delta^{\frac{1-2b}{2}} \left\| (|v_\delta|^2 - w^\varepsilon * |v_\delta|^2) v_\delta \right\|_{X^{0,b-1}} + C \delta^{\frac{1-2b}{2}} \left\| \left\{ w^\varepsilon * (|v_\delta|^2 - |v_\delta^\varepsilon|^2) \right\} v_\delta \right\|_{X^{0,b-1}} \\ &\quad + C \delta^{\frac{1-2b}{2}} \left\| \left\{ w^\varepsilon * |v_\delta^\varepsilon|^2 \right\} (v_\delta - v_\delta^\varepsilon) \right\|_{X^{0,b-1}}. \end{aligned} \quad (5.26)$$

Note that, in the above expressions, the quantity v_δ^ε is obtained from v^ε according to (5.12). We now estimate each of the terms on the right-hand side of (5.26) separately.

For the first term, we note that for fixed $x \in \Lambda$ we have

$$\left| |v_\delta(x)|^2 - (w^\varepsilon * |v_\delta|^2)(x) \right| \leq \int dy w^\varepsilon(x-y) |v_\delta(x) - v_\delta(y)| (|v_\delta(x)| + |v_\delta(y)|). \quad (5.27)$$

Here, we used (1.30) by which we obtain that

$$\int dx w^\varepsilon(x) = 1. \quad (5.28)$$

Moreover, we used the elementary inequality

$$||a_1|^2 - |a_2|^2| \leq |a_1 - a_2|(|a_1| + |a_2|). \quad (5.29)$$

We recall (5.4) and use Lemma 5.3 (vi) to note that

$$\begin{aligned} \left\| (|v_\delta|^2 - w^\varepsilon * |v_\delta|^2) v_\delta \right\|_{X^{0,b-1}} &\leq \left\| (|v_\delta|^2 - w^\varepsilon * |v_\delta|^2) v_\delta \right\|_{X^{0,-3/8}} \\ &\leq C \left\| (|v_\delta|^2 - w^\varepsilon * |v_\delta|^2) v_\delta \right\|_{L_{t,x}^{4/3}}, \end{aligned} \quad (5.30)$$

which by (5.27) is

$$\leq C \left\| w^\varepsilon(x-y) |v_\delta(x) - v_\delta(y)| |v_\delta(x)| \right\|_{L_t^{4/3} L_x^{4/3} L_y^1} + C \left\| w^\varepsilon(x-y) |v_\delta(x) - v_\delta(y)| |v_\delta(y)| \right\|_{L_t^{4/3} L_x^{4/3} L_y^1}. \quad (5.31)$$

Note that, in order to apply (5.27) in (5.30), it is crucial to use that we are estimating the $L_{t,x}^{4/3}$ norm and not the $X^{0,b-1}$ norm.

By Hölder's inequality in mixed-norm spaces and by the construction of v_δ in (5.12), the expression in (5.31) is

$$\begin{aligned} &\leq \left\| [x-y]^{s+\frac{1}{2}} w^\varepsilon(x-y) \right\|_{L_x^\infty L_y^2} \left\| \psi_\delta \right\|_{L_t^\infty} \left\| \frac{v(x) - v(y)}{[x-y]^{s+\frac{1}{2}}} \right\|_{L_t^\infty L_{x,y}^2} \left\| \psi_\delta \right\|_{L_t^{4/3}} \left\| v(x) \right\|_{L_t^\infty L_x^4} \\ &\quad + \left\| [x-y]^{s+\frac{1}{2}} w^\varepsilon(x-y) \right\|_{L_{x,y}^4} \left\| \psi_\delta \right\|_{L_t^\infty} \left\| \frac{v(x) - v(y)}{[x-y]^{s+\frac{1}{2}}} \right\|_{L_t^\infty L_{x,y}^2} \left\| \psi_\delta \right\|_{L_t^{4/3}} \left\| v(y) \right\|_{L_t^\infty L_y^4}. \end{aligned} \quad (5.32)$$

We now estimate (5.32). By (1.30) and by the assumption that $w \in C_c^\infty(\mathbb{R})$, we have that for all $1 \leq p < \infty$

$$\begin{aligned} \left\| [z]^{s+\frac{1}{2}} w^\varepsilon(z) \right\|_{L_z^p} &= \frac{1}{\varepsilon} \left(\int_0^1 dz z^{(s+\frac{1}{2})p} \left| w\left(\frac{z}{\varepsilon}\right) \right|^p \right)^{1/p} = \varepsilon^{s+\frac{1}{p}-\frac{1}{2}} \left(\int_0^{1/\varepsilon} d\tilde{z} \tilde{z}^{(s+\frac{1}{2})p} |w(\tilde{z})|^p \right)^{1/p} \\ &\leq C(s,p,w) \varepsilon^{s+\frac{1}{p}-\frac{1}{2}}. \end{aligned} \quad (5.33)$$

Here we used the change of variables $\tilde{z} = z/\varepsilon$. We now apply (5.33) with $p = 2$ and $p = 4$ and deduce that

$$\left\| [x-y]^{s+\frac{1}{2}} w^\varepsilon(x-y) \right\|_{L_x^\infty L_y^2} \leq C\varepsilon^s, \quad \left\| [x-y]^{s+\frac{1}{2}} w^\varepsilon(x-y) \right\|_{L_{x,y}^4} \leq C\varepsilon^{s-\frac{1}{4}}. \quad (5.34)$$

For the second inequality in (5.34), we also used the compactness of Λ . Furthermore, by Lemma 5.4, Lemma 5.3 (i) and since $v \in \Gamma$ for the set Γ defined as in (5.17), it follows that

$$\left\| \frac{v(x) - v(y)}{[x-y]^{s+\frac{1}{2}}} \right\|_{L_t^\infty L_{x,y}^2} \leq C \|v\|_{L_t^\infty H_x^s} \leq C \|v\|_{X^{s,b}} \leq C\delta^{\frac{1-2b}{2}} \|\phi_0\|_{H^s}. \quad (5.35)$$

Moreover, we note that, by Hölder's inequality, Sobolev embedding with $s \geq \frac{1}{4}$ and the same arguments as in (5.35), we have

$$\|v\|_{L_t^\infty L_x^4} \leq C \|v\|_{L_t^\infty H_x^s} \leq C\delta^{\frac{1-2b}{2}} \|\phi_0\|_{H^s}. \quad (5.36)$$

We use (5.34)–(5.36), as well as (5.9)–(5.10) to deduce that the expression in (5.32) is

$$\leq C \delta^{\frac{3}{4}+(1-2b)} \|\phi_0\|_{H^s}^2 \varepsilon^{s-1/4}. \quad (5.37)$$

We now estimate the second term on the right-hand of (5.26). By applying Lemma 5.3 (vi) as in (5.30), it follows that

$$\begin{aligned} \left\| \left\{ w^\varepsilon * (|v_\delta|^2 - |v_\delta^\varepsilon|^2) \right\} v_\delta \right\|_{X^{0,b-1}} &\leq C \left\| \left\{ w^\varepsilon * (|v_\delta|^2 - |v_\delta^\varepsilon|^2) \right\} v_\delta \right\|_{L_{t,x}^{4/3}} \\ &\leq C \left\| \left\{ w^\varepsilon * (|v_\delta - v_\delta^\varepsilon| (|v_\delta| + |v_\delta^\varepsilon|)) \right\} v_\delta \right\|_{L_{t,x}^{4/3}}. \end{aligned}$$

In the last inequality, we also used (5.29). By Hölder's and Young's inequality, it follows that this expression is

$$\begin{aligned} &\leq \|v_\delta - v_\delta^\varepsilon\|_{L_{t,x}^2} \left(\|v_\delta\|_{L_{t,x}^8} + \|v_\delta^\varepsilon\|_{L_{t,x}^8} \right) \|v_\delta\|_{L_{t,x}^8} \\ &\leq C \delta^{\frac{3}{4}} \|v - v^\varepsilon\|_{L_t^\infty L_x^2} \left(\|v\|_{L_t^\infty L_x^8} + \|v^\varepsilon\|_{L_t^\infty L_x^8} \right) \|v\|_{L_t^\infty L_x^8} \\ &\leq C \delta^{\frac{3}{4}} \|v - v^\varepsilon\|_{X^{0,b}} \left(\|v\|_{X^{s,b}} + \|v^\varepsilon\|_{X^{s,b}} \right) \|v\|_{X^{s,b}} \leq C \delta^{\frac{3}{4}+(1-2b)} \|\phi_0\|_{H^s}^2 \|v - v^\varepsilon\|_{X^{0,b}}. \quad (5.38) \end{aligned}$$

Above we used Sobolev embedding with $s \geq \frac{3}{8}$, Lemma 5.3 (i), the construction of $v, v_\delta, v_\delta^\varepsilon, v, v^\varepsilon$, as well as $\|w^\varepsilon\|_{L^1} = 1$, which follows from (5.28) since $w^\varepsilon \geq 0$.

The third term on the right-hand side of (5.26) is estimated in a similar way. Arguing as in (5.30), we need to estimate

$$\begin{aligned} \left\| \left\{ w^\varepsilon * |v_\delta^\varepsilon|^2 \right\} (v_\delta - v_\delta^\varepsilon) \right\|_{L_{t,x}^{4/3}} &\leq \|v_\delta^\varepsilon\|_{L_{t,x}^8}^2 \|v_\delta - v_\delta^\varepsilon\|_{L_{t,x}^2} \leq C \delta^{\frac{3}{4}} \|v^\varepsilon\|_{L_t^\infty L_x^8}^2 \|v - v^\varepsilon\|_{L_t^\infty L_x^2} \\ &\leq C \delta^{\frac{3}{4}} \|v^\varepsilon\|_{X^{s,b}}^2 \|v - v^\varepsilon\|_{X^{0,b}} \leq C \delta^{\frac{3}{4}+(1-2b)} \|\phi_0\|_{H^s}^2 \|v - v^\varepsilon\|_{X^{0,b}}. \quad (5.39) \end{aligned}$$

Here, we again used Hölder's inequality, Young's inequality, Sobolev embedding with $s \geq \frac{3}{8}$, Lemma 5.3 (i) and the construction of $v_\delta, v_\delta^\varepsilon, v^\varepsilon, w^\varepsilon$. Substituting (5.37)–(5.39) into (5.26), it follows that

$$\|v - v^\varepsilon\|_{X^{0,b}} \leq C \delta^{\theta_1} \|\phi_0\|_{H^s}^2 \varepsilon^{s-\frac{1}{4}} + C \delta^{\theta_1} \|\phi_0\|_{H^s}^2 \|v - v^\varepsilon\|_{X^{0,b}}, \quad (5.40)$$

where

$$\theta_1 := \frac{3}{4} + \frac{3(1-2b)}{2} > 0.$$

In particular, if we choose $\delta \equiv \delta(\|\phi_0\|_{H^s}) > 0$ possibly smaller than before so that the coefficient of $\|v - v^\varepsilon\|_{X^{0,b}}$ on the right-hand side of (5.40) is smaller than 1/2, it follows that

$$\|v - v^\varepsilon\|_{X^{0,b}} \leq C(\|\phi_0\|_{H^s}) \varepsilon^{s-\frac{1}{4}}. \quad (5.41)$$

By analogous arguments, we obtain more generally that for all $n \in \mathbb{N}$ we have

$$\begin{aligned} \|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}} &\leq C \delta^{\frac{1-2b}{2}} \|v_{(n)}(n\delta) - v_{(n)}^\varepsilon(n\delta)\|_{\mathfrak{H}} + C \delta^{\theta_1} \|v_{(n)}(n\delta)\|_{H^s}^2 \varepsilon^{s-\frac{1}{4}} \\ &\quad + C \delta^{\theta_1} \left(\|v_{(n)}(n\delta)\|_{H^s} + \|v_{(n)}^\varepsilon(n\delta)\|_{H^s} \right)^2 \|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}}. \quad (5.42) \end{aligned}$$

Note that the first term on the right-hand side of (5.42) appears because in general we consider different initial data $v_{(n)}(n\delta)$ and $v_{(n)}^\varepsilon(n\delta)$. We hence obtain the corresponding term on the right-hand side of (5.42) by Lemma 5.3 (ii).

In particular, if $1 \leq n \leq (T-1)/\delta$, we obtain by Lemma 5.3 (i), (5.24)–(5.25) and (5.42) that

$$\begin{aligned} \|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}} &\leq C \delta^{\frac{1-2b}{2}} \|v_{(n-1)} - v_{(n-1)}^\varepsilon\|_{X^{0,b}} \\ &\quad + C_1(\|\phi_0\|_{H^s}, T) \varepsilon^{s-\frac{1}{4}} + C_2(\|\phi_0\|_{H^s}, T) \delta^{\theta_1} \|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}}. \end{aligned}$$

Here we also assume that $\delta < 1$. In particular, choosing $\delta \equiv \delta(\|\phi_0\|_{H^s}, T) > 0$ even smaller than before such that $C_2(\|\phi_0\|_{H^s}, T) \delta^{\theta_1} < 1/2$, it follows that

$$\|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}} \leq C(\|\phi_0\|_{H^s}, T) \|v_{(n-1)} - v_{(n-1)}^\varepsilon\|_{X^{0,b}} + C(\|\phi_0\|_{H^s}, T) \varepsilon^{s-\frac{1}{4}}, \quad (5.43)$$

for all $1 \leq n \leq (T-1)/\delta$. Indeed, by (5.25), we can take

$$\delta \equiv \delta \left(\sup_{[0,T]} \|u(t)\|_{H^s} + \sup_{\varepsilon > 0} \sup_{[0,T]} \|u^\varepsilon(t)\|_{H^s} \right) = \delta(\|\phi_0\|_{H^s}, T) > 0$$

in order to obtain (5.43).

Iterating (5.43) and recalling (5.41), it follows that for all $0 \leq n \leq (T-1)/\delta$ we have

$$\|v_{(n)} - v_{(n)}^\varepsilon\|_{X^{0,b}} \leq C(\|\phi_0\|_{H^s}, T) \varepsilon^{s-\frac{1}{4}}. \quad (5.44)$$

Using Lemma 5.3 (i), (5.24) and (5.44), it follows that

$$\|u - u^\varepsilon\|_{L_{[0,T]}^\infty \mathfrak{H}} \leq C(\|\phi_0\|_{H^s}, T) \varepsilon^{s-\frac{1}{4}},$$

from where we deduce the claim since $s \geq \frac{3}{8}$. \square

Before proceeding to the proof of Theorem 1.6 we record the following elementary lemma.

Lemma 5.5. *Let $(Z_k)_{k \in \mathbb{N}}$ be an increasing family of sets (i.e. $Z_k \subset Z_{k+1}$), and set $Z := \bigcup_{k \in \mathbb{N}} Z_k$. For $\varepsilon, \tau > 0$ let $f, f^\varepsilon, f_\tau^\varepsilon : Z \rightarrow \mathbb{C}$ be functions which satisfy the following properties.*

- (i) *For each fixed $k \in \mathbb{N}$ and $\varepsilon > 0$ we have $\lim_{\tau \rightarrow \infty} f_\tau^\varepsilon(\zeta) = f^\varepsilon(\zeta)$ uniformly in $\zeta \in Z_k$.*
- (ii) *For each fixed $k \in \mathbb{N}$ we have $\lim_{\varepsilon \rightarrow 0} f^\varepsilon(\zeta) = f(\zeta)$ uniformly in $\zeta \in Z_k$.*

Then there exists a sequence of positive numbers (ε_τ) , with $\lim_{\tau \rightarrow \infty} \varepsilon_\tau = 0$, such that

$$\lim_{\tau \rightarrow \infty} f_\tau^{\varepsilon_\tau}(\zeta) = f(\zeta),$$

for all $\zeta \in Z$.

Proof. Assumptions (i) and (ii) combined with a diagonal argument imply that, for a fixed $k \in \mathbb{N}$, there exists a sequence of positive numbers (ε_τ^k) , with $\lim_{\tau \rightarrow \infty} \varepsilon_\tau^k = 0$, such that $\lim_{\tau \rightarrow \infty} f_\tau^{\varepsilon_\tau^k}(\zeta) = f(\zeta)$, uniformly in $\zeta \in Z_k$. Using a further diagonal argument, we extract a diagonal sequence (ε_τ) from (ε_τ^k) such that $\lim_{\tau \rightarrow \infty} f_\tau^{\varepsilon_\tau}(\zeta) = f(\zeta)$, for all $\zeta \in Z$. \square

Proof of Theorem 1.6. We shall apply Lemma 5.5 for the following choices of the sets Z_k and Z

$$Z := \left\{ (m, t_1, \dots, t_m, p_1, \dots, p_m, \xi^1, \dots, \xi^m) : m \in \mathbb{N}, t_i \in \mathbb{R}, p_i \in \mathbb{N}, \xi^i \in \mathcal{L}(\mathfrak{H}^{(p_i)}) \right\}$$

$$Z_k := \left\{ (m, t_1, \dots, t_m, p_1, \dots, p_m, \xi^1, \dots, \xi^m) \in Z : m \leq k, |t_i| \leq k, p_i \leq k, \|\xi^i\| \leq k \right\}$$

and the functions

$$f_{\sharp}^{\varepsilon}(\zeta) := \rho_{\sharp}^{\varepsilon}(\Psi_{\sharp}^{t_1, \varepsilon} \Theta_{\sharp}(\xi^1) \dots \Psi_{\sharp}^{t_m, \varepsilon} \Theta_{\sharp}(\xi^m)), \quad f(\zeta) := \rho(\Psi^{t_1} \Theta(\xi^1) \dots \Psi^{t_m} \Theta(\xi^m)),$$

where \sharp stands for either nothing or τ .

By Theorem 1.4, here used for the choice $w = w^{\varepsilon}$, Lemma 5.5, and Remark 4.1, it suffices to show that, for fixed $k \in \mathbb{N}$,

$$\lim_{\varepsilon \rightarrow 0} \rho^{\varepsilon}(\Psi^{t_1, \varepsilon} \Theta(\xi^1) \dots \Psi^{t_m, \varepsilon} \Theta(\xi^m)) = \rho(\Psi^{t_1} \Theta(\xi^1) \dots \Psi^{t_m} \Theta(\xi^m)), \quad (5.45)$$

uniformly in the parameters

$$m \leq k, |t_i| \leq k, p_i \leq k, \|\xi^i\| \leq k, i = 1, \dots, m. \quad (5.46)$$

In the sequel, we set $w = w^{\varepsilon}$ and we add a superscript ε to any quantity defined in terms of w to indicate that in its definition w is replaced by w^{ε} . For instance, we write the expectation $\rho^{\varepsilon}(X)$ as in (1.27), the classical interaction $\mathcal{W}^{\varepsilon}$ defined as in (1.24) and $\Psi^{t, \varepsilon} \Theta(\xi)$ given as in Definition 1.3, all with this modification. In addition, we write the classical interaction \mathcal{W} defined as in (1.24) with w formally set to equal the delta function. With these conventions we define

$$\tilde{\rho}_z^{\varepsilon}(X) := \int X e^{-z\mathcal{W}^{\varepsilon}} d\mu, \quad \tilde{\rho}_z(X) := \int X e^{-z\mathcal{W}} d\mu$$

for a random variable X and $\operatorname{Re} z \geq 0$. In particular, we have

$$\rho^{\varepsilon}(X) = \frac{\tilde{\rho}_1^{\varepsilon}(X)}{\tilde{\rho}_1^{\varepsilon}(1)}, \quad \rho(X) = \frac{\tilde{\rho}_1(X)}{\tilde{\rho}_1(1)}. \quad (5.47)$$

Let us first observe that

$$\lim_{\varepsilon \rightarrow 0} \mathcal{W}^{\varepsilon} = \mathcal{W} \quad \text{almost surely.} \quad (5.48)$$

Indeed, using (5.28)–(5.29), we obtain

$$|2(\mathcal{W} - \mathcal{W}^{\varepsilon})| \leq \int dx dy w^{\varepsilon}(x - y) |\phi(x) - \phi(y)| (|\phi(x)| + |\phi(y)|) |\phi(x)|^2.$$

Set $s = \frac{3}{8}$. Note that, since $s < \frac{1}{2}$, the free classical field ϕ defined in (1.22) is in $H^s(\Lambda)$ almost surely. We now apply Hölder's inequality in mixed norm spaces similarly as in (5.32) to deduce that this expression is

$$\begin{aligned} &\leq \left\| [x - y]^{s + \frac{1}{2}} w^{\varepsilon}(x - y) \right\|_{L_x^{\infty} L_y^2} \left\| \frac{\phi(x) - \phi(y)}{[x - y]^{s + \frac{1}{2}}} \right\|_{L_{x,y}^2} \|\phi(x)\|_{L_x^6}^3 \\ &+ \left\| [x - y]^{s + \frac{1}{2}} w^{\varepsilon}(x - y) \right\|_{L_x^{\infty} L_y^4} \left\| \frac{\phi(x) - \phi(y)}{[x - y]^{s + \frac{1}{2}}} \right\|_{L_{x,y}^2} \|\phi(y)\|_{L_y^4} \|\phi(x)\|_{L_x^4}^2 \leq C \varepsilon^{s - \frac{1}{4}} \|\phi\|_{H^s}^4. \end{aligned} \quad (5.49)$$

Here we used (5.34), Lemma 5.4 and Sobolev embedding with $s \geq \frac{1}{4}$. The claim (5.48) now follows from (5.49) since $\phi \in H^s(\Lambda)$ almost surely.

Since $\mathcal{W}^\varepsilon, \mathcal{W} \geq 0$, it follows from (5.48) and the dominated convergence theorem that

$$\lim_{\varepsilon \rightarrow 0} \tilde{\rho}_1^\varepsilon(1) = \tilde{\rho}_1(1). \quad (5.50)$$

In particular, by (5.47) and (5.50) we deduce that (5.45) is equivalent to showing that

$$\lim_{\varepsilon \rightarrow 0} \tilde{\rho}_1^\varepsilon \left(\Psi^{t_1, \varepsilon} \Theta(\xi^1) \dots \Psi^{t_m, \varepsilon} \Theta(\xi^m) \right) = \tilde{\rho}_1 \left(\Psi^{t_1} \Theta(\xi^1) \dots \Psi^{t_m} \Theta(\xi^m) \right), \quad (5.51)$$

uniformly in (5.46). In order to prove (5.51), we note that, by construction of $\Psi^{t, \varepsilon}, \Psi^t$ and Proposition 5.1, we have that, for $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$,

$$\lim_{\varepsilon \rightarrow 0} \Psi^{t, \varepsilon} \Theta(\xi) = \Psi^t \Theta(\xi) \quad \text{in } \mathfrak{H} \text{ almost surely.} \quad (5.52)$$

The convergence in (5.52) is uniform in $|t| \leq k, p_i \leq k, \|\xi^i\| \leq k$. Indeed, we write

$$\Psi^{t, \varepsilon} \Theta(\xi) = \left\langle (S_t^\varepsilon \phi)^{\otimes k}, \xi (S_t^\varepsilon \phi)^{\otimes k} \right\rangle_{\mathfrak{H}^{\otimes k}}, \quad \Psi^t \Theta(\xi) = \left\langle (S_t \phi)^{\otimes k}, \xi (S_t \phi)^{\otimes k} \right\rangle_{\mathfrak{H}^{\otimes k}},$$

where S_t^ε and S_t denote the flow maps of (5.2) and (5.1) respectively. We consider the initial data ϕ_0 given by the free classical field ϕ (1.22). Let us recall that $\phi \in H^s \subset \mathfrak{H}$ almost surely. Proposition 5.1 then implies that $\lim_{\varepsilon \rightarrow 0} (S_t^\varepsilon \phi)^{\otimes k} = (S_t \phi)^{\otimes k}$ in $\mathfrak{H}^{\otimes k}$ almost surely. We deduce (5.52) since $\xi \in \mathcal{L}(\mathfrak{H}^{(p)})$.

In particular, from (5.48) and (5.52) it follows that

$$\lim_{\varepsilon \rightarrow 0} \Psi^{t_1, \varepsilon} \Theta(\xi^1) \dots \Psi^{t_m, \varepsilon} \Theta(\xi^m) e^{-\mathcal{W}^\varepsilon} = \Psi^{t_1} \Theta(\xi^1) \dots \Psi^{t_m} \Theta(\xi^m) e^{-\mathcal{W}} \quad \text{almost surely.} \quad (5.53)$$

Furthermore, by conservation of mass for (5.2) and since $\mathcal{W}^\varepsilon \geq 0$ by construction, it follows that for all $\varepsilon > 0$ we have

$$\left| \Psi^{t_1, \varepsilon} \Theta(\xi^1) \dots \Psi^{t_m, \varepsilon} \Theta(\xi^m) e^{-\mathcal{W}^\varepsilon} \right| \leq \|\xi^1\| \dots \|\xi^m\| \|\phi\|_{\mathfrak{H}}^{2(p_1 + \dots + p_m)} \in L^1(d\mu). \quad (5.54)$$

We now deduce (5.51) from (5.53)–(5.54) and the dominated convergence theorem. \square

A. $X^{\sigma, b}$ spaces: proof of Lemma 5.3.

In this appendix we present the proof of Lemma 5.3. We emphasize that this is done for the convenience of the reader and that it is not an original contribution of the paper.

Proof of Lemma 5.3. We recall the definition of b given in (5.4).

We first prove part (i). The proof is analogous to the proof the Sobolev embedding $H_t^b \hookrightarrow L_t^\infty$. We use the Fourier inversion formula in the time variable and write

$$\hat{f}(k, t) = \int_{-\infty}^{\infty} d\eta \tilde{f}(k, \eta) e^{2\pi i \eta t}. \quad (A.1)$$

In (A.1),

$$\hat{f}(k, t) = \int_{\Lambda} dx f(x, t) e^{-2\pi i k x}$$

denotes the Fourier transform in the space variable. In particular, using the Cauchy-Schwarz inequality in η in (A.1) and recalling that $b > 1/2$, it follows that

$$|\hat{f}(k, t)| \leq C(b) \left(\int_{-\infty}^{\infty} d\eta |\tilde{f}(k, \eta)|^2 (1 + |\eta + 2\pi k^2|)^{2b} \right)^{1/2}. \quad (\text{A.2})$$

Claim (i) follows from (A.2) and Definition 5.2.

Claims analogous to (ii)-(iv) were proved for $X^{\sigma, b}$ corresponding to the Airy equation in the non-periodic setting [15, Lemmas 3.1–3.3]. The bounds for the Schrödinger equation follow in the same way, since we are estimating integrals in the Fourier variable η dual to time. For completeness, we give the proofs of (ii)-(iv).

We proceed with the proof of (ii). By density, it suffices to consider $\Phi \in \mathcal{S}(\Lambda_x)$. Let us note that, for fixed $x \in \Lambda$

$$\psi(t/\delta) e^{it\Delta} \Phi = \psi(t/\delta) \sum_k e^{2\pi i k x - 4\pi^2 i k^2 t} \hat{\Phi}(k),$$

from where we deduce that

$$(\psi(t/\delta) e^{it\Delta} \Phi)^\sim(k, \tau) = \delta \hat{\psi}(\delta(\eta + 2\pi k^2)) \hat{\Phi}(k).$$

Hence

$$\begin{aligned} \|\psi(t/\delta) e^{it\Delta} \Phi\|_{X^{\sigma, b}}^2 &= \sum_k (1 + |2\pi k|)^{2\sigma} |\hat{\Phi}(k)|^2 \left[\delta^2 \int_{-\infty}^{\infty} d\eta \left| \hat{\psi}(\delta(\eta + 2\pi k^2)) \right|^2 (1 + |\eta + 2\pi k^2|)^{2b} \right] \\ &= \|\Phi\|_{H^\sigma}^2 \left[\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}(\delta\eta)|^2 (1 + |\eta|)^{2b} \right]. \end{aligned} \quad (\text{A.3})$$

By scaling we obtain that the following estimates hold.

$$\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}(\delta\eta)|^2 \leq C(\psi). \quad (\text{A.4})$$

$$\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}(\delta\eta)|^2 |\eta|^{2b} \leq C(b, \psi) \delta^{1-2b}. \quad (\text{A.5})$$

Claim (ii) follows by substituting (A.4)–(A.5) into (A.3) and using

$$(1 + |\eta|)^{2b} \leq C(b)(1 + |\eta|^{2b}).$$

(In the sequel, we use the latter elementary inequality repeatedly without explicit mention).

We now prove (iii). Let us note that

$$(\psi(t/\delta)f)^\sim(k, \eta) = \tilde{f}(k, \eta) *_{\eta} (\delta \hat{\psi}(\delta \cdot)) \quad (\text{A.6})$$

where $*_\eta$ denotes convolution in η . From (A.6) and Definition 5.2, it follows that (iii) is equivalent to showing that for all $h = h(t)$ and $a \in \mathbb{R}$ we have

$$\int_{-\infty}^{\infty} d\eta \left| \hat{h} *_\eta (\delta \hat{\psi}(\delta \cdot))(\eta) \right|^2 (1 + |\eta + a|)^{2b} \leq C(b, \psi) \delta^{1-2b} \int_{-\infty}^{\infty} d\eta |\hat{h}(\eta)|^2 (1 + |\eta + a|)^{2b}. \quad (\text{A.7})$$

By Young's inequality, it follows that

$$\int_{-\infty}^{\infty} d\eta \left| \hat{h} *_\eta (\delta \hat{\psi}(\delta \cdot))(\eta) \right|^2 \leq C(\psi) \int_{-\infty}^{\infty} d\eta |\hat{h}(\eta)|^2. \quad (\text{A.8})$$

Moreover, we write

$$\begin{aligned} \int_{-\infty}^{\infty} d\eta \left| \hat{h} *_\eta (\delta \hat{\psi}(\delta \cdot))(\eta) \right|^2 |\eta + a|^{2b} &= \int_{-\infty}^{\infty} d\eta \left| \hat{h} *_\eta (\delta \hat{\psi}(\delta \cdot))(\eta - a) \right|^2 |\eta|^{2b} \\ &= C(b) \int_{-\infty}^{\infty} dt \left| |\partial|^b (e^{2\pi i a t} h(t) \psi(\delta^{-1} t)) \right|^2 = C(b) \left\| |\partial|^b (e^{2\pi i a t} h \psi(\delta^{-1} \cdot)) \right\|_{L_t^2}^2. \end{aligned} \quad (\text{A.9})$$

Here we use the notation $|\partial|^b$ for the fractional differentiation operator given by

$$(|\partial|^b g)^\wedge(\eta) = |2\pi\eta|^b \hat{g}(\eta).$$

We now refer to the result of [14, Theorem A.12] (c.f. also [15, Theorem 2.8]) which states that for all $\alpha \in (0, 1)$ and $p \in (1, \infty)$ we have

$$\left\| |\partial|^\alpha (fg) - f |\partial|^\alpha g \right\|_{L^p} \leq C(\alpha, p) \|g\|_{L^\infty} \left\| |\partial|^\alpha f \right\|_{L^p}. \quad (\text{A.10})$$

Taking $\alpha = b$, $p = 2$, $f = e^{2\pi i a t} h$ and $g = \psi(\delta^{-1} \cdot)$ in (A.10) we obtain

$$\begin{aligned} \left\| |\partial|^b (e^{2\pi i a t} h \psi(\delta^{-1} \cdot)) - e^{2\pi i a t} h |\partial|^b (\psi(\delta^{-1} \cdot)) \right\|_{L_t^2} &\leq C(b) \|\psi(\delta^{-1} \cdot)\|_{L_t^\infty} \left\| |\partial|^b (e^{2\pi i a t} h) \right\|_{L_t^2} \\ &\leq C(b, \psi) \left\| |\partial|^b (e^{2\pi i a t} h) \right\|_{L_t^2}. \end{aligned} \quad (\text{A.11})$$

By Plancherel's theorem we have

$$\left\| |\partial|^b (e^{2\pi i a t} h) \right\|_{L_t^2}^2 = C(b) \int_{-\infty}^{\infty} d\eta |\hat{h}(\eta - a)|^2 |\eta|^{2b} = C(b) \int_{-\infty}^{\infty} d\eta |\hat{h}(\eta)|^2 |\eta + a|^{2b}. \quad (\text{A.12})$$

From (A.8)–(A.9) and (A.11)–(A.12), we deduce that (A.7) follows if we show that

$$I := \left\| e^{2\pi i a t} h |\partial|^b (\psi(\delta^{-1} \cdot)) \right\|_{L_t^2} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \left(\int_{-\infty}^{\infty} d\eta |\hat{h}(\eta)|^2 (1 + |\eta + a|)^{2b} \right)^{1/2}. \quad (\text{A.13})$$

Applying Hölder's inequality and Sobolev embedding with $b > \frac{1}{2}$, it follows that

$$\begin{aligned} I &\leq \|e^{2\pi i a t} h\|_{L_t^\infty} \left\| |\partial|^b (\psi(\delta^{-1} \cdot)) \right\|_{L_t^2} \leq C(b) \|e^{2\pi i a t} h\|_{H_t^b} \left\| |\partial|^b (\psi(\delta^{-1} \cdot)) \right\|_{L_t^2} \\ &= C(b) \left(\int_{-\infty}^{\infty} d\eta |\hat{h}(\eta)|^2 (1 + |\eta + a|)^{2b} \right)^{1/2} \left\| |\partial|^b (\psi(\delta^{-1} \cdot)) \right\|_{L_t^2}. \end{aligned} \quad (\text{A.14})$$

By scaling, we compute

$$\left\| |\partial|^b \left(\psi(\delta^{-1} \cdot) \right) \right\|_{L_t^2} = C(b) \delta^{\frac{1-2b}{2}} \left(\int_{-\infty}^{\infty} d\eta |\eta|^{2b} |\hat{\psi}(\eta)|^2 \right)^{1/2} \leq C(b, \psi) \delta^{\frac{1-2b}{2}}. \quad (\text{A.15})$$

By (A.14)–(A.15), we deduce (A.13), which in turn implies (A.7). The claim (iii) now follows.

We now prove (iv). By density, it suffices to consider $f \in \mathcal{S}(\Lambda_x \times \mathbb{R}_t)$. We write

$$J := \psi(t/\delta) \int_0^t dt' e^{i(t-t')\Delta} f(t') = \psi(t/\delta) \int_0^t dt' \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} e^{-4\pi^2 i k^2 t} e^{2\pi i \eta t' (\eta + 2\pi k^2)}. \quad (\text{A.16})$$

By the assumptions on f we can interchange the orders of integration so that we first integrate in t' . Evaluating the t' integral, it follows that

$$J = \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \frac{e^{2\pi i \eta t} - e^{-4\pi^2 i k^2 t}}{2\pi i (\eta + 2\pi k^2)} = I_1 + I_2, \quad (\text{A.17})$$

where

$$I_1 := \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \Xi(\eta + 2\pi k^2) \frac{e^{2\pi i \eta t} - e^{-4\pi^2 i k^2 t}}{2\pi i (\eta + 2\pi k^2)} \quad (\text{A.18})$$

$$I_2 := \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \left(1 - \Xi(\eta + 2\pi k^2) \right) \frac{e^{2\pi i \eta t} - e^{-4\pi^2 i k^2 t}}{2\pi i (\eta + 2\pi k^2)}, \quad (\text{A.19})$$

for a function $\Xi \in C_c^\infty(\mathbb{R})$ such that

$$\Xi = 1 \quad \text{for } |y| \leq 1/2, \quad \text{and } \Xi = 0 \quad \text{for } |y| > 1. \quad (\text{A.20})$$

We first consider I_1 . By writing a Taylor expansion for the factor

$$\frac{e^{2\pi i \eta t} - e^{-4\pi^2 i k^2 t}}{2\pi i (\eta + 2\pi k^2)} = e^{-4\pi^2 i k^2 t} \frac{e^{2\pi i t(\eta + 2\pi k^2)} - 1}{2\pi i (\eta + 2\pi k^2)}$$

in the integrand of (A.18), we have

$$I_1 = \sum_{l=1}^{\infty} \frac{(2\pi i)^{l-1} t^l}{l!} \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \Xi(\eta + 2\pi k^2) e^{-4\pi^2 i k^2 t} (\eta + 2\pi k^2)^{l-1}. \quad (\text{A.21})$$

Here, we can justify taking the sum in l outside of the integral in η and the sum in k using the assumption that $f \in \mathcal{S}(\Lambda_x \times \mathbb{R}_t)$ as before. Setting

$$\psi_l(y) := y^l \psi(y) \quad (\text{A.22})$$

and using the Fourier representation of $e^{it\Delta}$, we can rewrite (A.21) as

$$I_1 = \sum_{l=1}^{\infty} \frac{(2\pi i)^{l-1} \delta^l}{l!} \psi_l(t/\delta) \left[e^{it\Delta} \left(\int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \Xi(\eta + 2\pi k^2) (\eta + 2\pi k^2)^{l-1} \right) \right]. \quad (\text{A.23})$$

Using the triangle inequality and (A.3), we obtain from (A.23) that

$$\begin{aligned} \|I_1\|_{X^{\sigma,b}} &\leq \sum_{l=1}^{\infty} \left(\frac{(2\pi)^{l-1} \delta^l}{l!} \left[\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}_l(\delta\eta)|^2 (1+|\eta|)^{2b} \right]^{1/2} \right. \\ &\quad \left. \times \left\| \left(\int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \Xi(\eta + 2\pi k^2) (\eta + 2\pi k^2)^{l-1} \right) \right\|_{H^\sigma} \right). \end{aligned} \quad (\text{A.24})$$

By scaling, we compute

$$\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}_l(\delta\eta)|^2 = \delta \|\psi_l\|_{L^2}^2. \quad (\text{A.25})$$

$$\delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}_l(\delta\eta)|^2 |\eta|^{2b} = \delta^{1-2b} \|\psi_l\|_{\dot{H}^b}^2. \quad (\text{A.26})$$

In particular, from (A.25)–(A.26), we deduce that

$$\begin{aligned} \delta^2 \int_{-\infty}^{\infty} d\eta |\hat{\psi}_l(\delta\eta)|^2 (1+|\eta|)^{2b} &\leq C(b) \delta^{1-2b} \left(\|\psi_l\|_{L^2}^2 + \|\psi_l\|_{\dot{H}^b}^2 \right) \leq C(b) \delta^{1-2b} \left(\|\psi_l\|_{L^2}^2 + \|\psi_l\|_{\dot{H}^1}^2 \right) \\ &\leq C(b) \delta^{1-2b} \left(\|y^l \psi\|_{L^2}^2 + \|y^l \psi'\|_{L^2}^2 + \|ly^{l-1} \psi\|_{L^2}^2 \right) \leq C(b) \delta^{1-2b} (C(\psi))^l. \end{aligned} \quad (\text{A.27})$$

Above we used (A.22) and the assumption that $\psi \in C_c^\infty(\mathbb{R})$.

Moreover, we have

$$\begin{aligned} &\left\| \left(\int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \Xi(\eta + 2\pi k^2) (\eta + 2\pi k^2)^{l-1} \right) \right\|_{H^\sigma}^2 \\ &= \sum_k (1 + |2\pi k|)^{2\sigma} \left| \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) \Xi(\eta + 2\pi k^2) (\eta + 2\pi k^2)^{l-1} \right|^2 \\ &\leq C(\Xi) \sum_k (1 + |2\pi k|)^{2\sigma} \left(\int_{|\eta+2\pi k^2| \leq 1} d\eta |\tilde{f}(k, \eta)| \right)^2, \end{aligned}$$

uniformly in l . In the above line, we used the assumptions (A.20) on the function Ξ . In particular, the above expression is

$$\begin{aligned} &\leq C(\Xi) \sum_k (1 + |2\pi k|)^{2\sigma} \left(\int_{|\eta+2\pi k^2| \leq 1} d\eta \frac{|\tilde{f}(k, \eta)|}{1 + |\eta + 2\pi k^2|} \right)^2 \\ &\leq C(\Xi) \sum_k (1 + |2\pi k|)^{2\sigma} \left(\int_{-\infty}^{\infty} d\eta \frac{|\tilde{f}(k, \eta)|}{(1 + |\eta + 2\pi k^2|)^{1-b}} \frac{1}{(1 + |\eta + 2\pi k^2|)^b} \right)^2 \\ &\leq C(b, \Xi) \left(\int_{-\infty}^{\infty} d\eta \sum_k (1 + |2\pi k|)^{2\sigma} (1 + |\eta + 2\pi k^2|)^{2(b-1)} |\tilde{f}(k, \eta)|^2 \right) = C(b, \Xi) \|f\|_{X^{s,b-1}}^2. \end{aligned} \quad (\text{A.28})$$

In the last inequality we used the Cauchy-Schwarz inequality in η and the assumption that $b > \frac{1}{2}$. Combining (A.24), (A.27)–(A.28), it follows that

$$\|I_1\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b-1}}. \quad (\text{A.29})$$

(Since Ξ is chosen as an arbitrary C_c^∞ function satisfying (A.20), we do not keep track of the dependence of the implied constant on this function).

We now consider I_2 as defined in (A.19). We write

$$I_2 = I_{2,1} - I_{2,2}, \quad (\text{A.30})$$

where

$$\begin{aligned} I_{2,1} &:= \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \left(1 - \Xi(\eta + 2\pi k^2)\right) \frac{e^{2\pi i \eta t}}{2\pi i(\eta + 2\pi k^2)}. \\ I_{2,2} &:= \psi(t/\delta) \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) e^{2\pi i k x} \left(1 - \Xi(\eta + 2\pi k^2)\right) \frac{e^{-4\pi^2 i k^2 t}}{2\pi i(\eta + 2\pi k^2)}. \end{aligned}$$

We first estimate $I_{2,1}$. By claim (iii), we have

$$\|I_{2,1}\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \left\| \int_{-\infty}^{\infty} d\eta \sum_k \tilde{f}(k, \eta) \frac{(1 - \Xi(\eta + 2\pi k^2))}{\eta + 2\pi k^2} e^{2\pi i k x + 2\pi i \eta t} \right\|_{X^{\sigma,b}}. \quad (\text{A.31})$$

Note that, by the assumptions (A.20) on Ξ we have

$$\left| \tilde{f}(k, \eta) \frac{(1 - \Xi(\eta + 2\pi k^2))}{\eta + 2\pi k^2} \right| \leq C(\Xi) \frac{|\tilde{f}(k, \eta)|}{(1 + |\eta + 2\pi k^2|)}. \quad (\text{A.32})$$

Combining (A.31)–(A.32) and recalling the definition of $\|\cdot\|_{X^{\sigma,b}}$ we deduce that

$$\begin{aligned} \|I_{2,1}\|_{X^{\sigma,b}} &\leq C(b, \psi, \Xi) \delta^{\frac{1-2b}{2}} \left\| (1 + |2\pi k|)^\sigma (1 + |\eta + 2\pi k^2|)^{b-1} |\tilde{f}(k, \eta)| \right\|_{L_{\eta k}^2} \\ &= C(b, \psi, \Xi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b-1}}. \end{aligned} \quad (\text{A.33})$$

In particular, in the last line, we used that the $X^{\sigma,b}$ norm depends only on the absolute value of the spacetime Fourier transform.

We now estimate $I_{2,2}$. Let us note that

$$I_{2,2} = \psi(t/\delta) e^{it\Delta} \left[\sum_k \left(\int_{-\infty}^{\infty} d\eta \tilde{f}(k, \eta) \frac{(1 - \Xi(\eta + 2\pi k^2))}{2\pi i(\eta + 2\pi k^2)} \right) e^{2\pi i k x} \right]. \quad (\text{A.34})$$

From (A.34) and part (ii) we obtain

$$\|I_{2,2}\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \left\| \sum_k \left(\int_{-\infty}^{\infty} d\eta \tilde{f}(k, \eta) \frac{(1 - \Xi(\eta + 2\pi k^2))}{2\pi i(\eta + 2\pi k^2)} \right) e^{2\pi i k x} \right\|_{H^\sigma},$$

which, by using the definition of $\|\cdot\|_{H^\sigma}$ and (A.32) is

$$\leq C(b, \psi, \Xi) \delta^{\frac{1-2b}{2}} \left[\sum_k (1 + |2\pi k|)^{2\sigma} \left(\int_{-\infty}^{\infty} d\eta \frac{|\tilde{f}(k, \eta)|}{(1 + |\eta + 2\pi k^2|)} \right)^2 \right]^{1/2}. \quad (\text{A.35})$$

By writing

$$\frac{1}{(1 + |\eta + 2\pi k^2|)} = \frac{1}{(1 + |\eta + 2\pi k^2|)^{1-b}} \frac{1}{(1 + |\eta + 2\pi k^2|)^b}$$

in the integrand in (A.35) and applying the Cauchy-Schwarz inequality in η analogously as in the proof of (A.28) above, it follows that

$$\begin{aligned} \|I_{2,2}\|_{X^{\sigma,b}} &\leq C(b, \psi, \Xi) \delta^{\frac{1-2b}{2}} \left[\sum_k (1 + |2\pi k|)^{2\sigma} \int_{-\infty}^{\infty} d\eta \frac{|\tilde{f}(k, \eta)|^2}{(1 + |\eta + 2\pi k^2|)^{2(1-b)}} \right]^{1/2} \\ &= C(b, \psi, \Xi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b-1}}. \end{aligned} \quad (\text{A.36})$$

From (A.30), (A.33) and (A.36), we obtain

$$\|I_2\|_{X^{\sigma,b}} \leq C(b, \psi) \delta^{\frac{1-2b}{2}} \|f\|_{X^{\sigma,b-1}}. \quad (\text{A.37})$$

(As in (A.29), we do not emphasize the Ξ -dependence in the implied constant). Claim (iv) now follows from (A.16)–(A.17), (A.29) and (A.37).

Claim (v) is proved in [3, Proposition 2.6]. For an alternative proof, see also [27, Proposition 2.13]. Claim (vi) follows from part (v) by duality.

Finally, we prove claim (vii). By part (v), it suffices to prove

$$\|\psi(t/\delta)f\|_{X^{0,3/8}} \leq C(b) \delta^{\theta_0} \|f\|_{X^{0,b}} \quad (\text{A.38})$$

for appropriately chosen $\theta_0 > 0$. In order to prove claim (A.38) we first note that

$$X^{0, \frac{1}{4}+} \hookrightarrow L_t^4 L_x^2, \quad (\text{A.39})$$

which follows by using $X^{0,0} = L_{t,x}^2$, part (i) with $\sigma = 0$ and interpolation. By an additional interpolation step, it follows that

$$\|\psi(t/\delta)f\|_{X^{0,3/8}} \leq \|\psi(t/\delta)f\|_{X^{0,0}}^\theta \|\psi(t/\delta)f\|_{X^{0,b}}^{1-\theta} \quad \text{for } \theta := \frac{b - \frac{3}{8}}{b} = \frac{1}{4} + . \quad (\text{A.40})$$

We have, by Hölder's inequality and (A.39)

$$\|\psi(t/\delta)f\|_{X^{0,0}} = \|\psi(t/\delta)f\|_{L_{t,x}^2} \leq \|\psi(t/\delta)\|_{L_t^4} \|f\|_{L_t^4 L_x^2} \leq C\delta^{\frac{1}{4}} \|f\|_{X^{0, \frac{1}{4}+}} \leq C\delta^{\frac{1}{4}} \|f\|_{X^{0,b}}. \quad (\text{A.41})$$

From (A.40)–(A.41) and part (iii), we obtain claim (vii) with

$$\theta_0 := \frac{\theta}{4} + (1 - \theta) \frac{1 - 2b}{2} > 0. \quad (\text{A.42})$$

□

References

- [1] F. Benaych-Georges and A. Knowles. Lectures on the local semicircle law for Wigner matrices. *Panoramas et Synthèses*, SMF, Vol. **53** (2018).
- [2] A. Bényi, T. Oh. The Sobolev inequality on the torus revisited. *Publ. Math. Debrecen* **83** (2013), no. 3, 359–374.
- [3] J. Bourgain. Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, I: Schrödinger equations. *Geom. Funct. Anal.* **3** (1993), 107–156.
- [4] J. Bourgain. Periodic nonlinear Schrödinger equation and invariant measures. *Comm. Math. Phys.* **166** (1994), 1–26.
- [5] J. Bourgain. On the Cauchy problem and invariant measure problem for the periodic Zakharov system. *Duke Math. J.* **76** (1994), 175–202.
- [6] J. Bourgain. Invariant measures for the 2D-defocusing nonlinear Schrödinger equation. *Comm. Math. Phys.* **176** (1996), 421–445.
- [7] J. Bourgain. Invariant measures for the Gross-Pitaevskii equation. *J. Math. Pures Appl.* **76** (1997), 649–702.
- [8] J. Bourgain. Invariant measures for NLS in infinite volume. *Comm. Math. Phys.* **210** (2000), 605–620.
- [9] T. Cazenave. Semilinear Schrödinger Equations. *AMS, CIMS Lecture Notes*, 2003.
- [10] J. Fröhlich, A. Knowles and A. Pizzo. Atomism and quantization. *J. Phys. A* **40** (2007), 3033–3045.
- [11] J. Fröhlich, A. Knowles, B. Schlein and V. Sohinger. Gibbs measures of nonlinear Schrödinger equations as limits of many-body quantum states in dimensions $d \leq 3$. *Comm. Math. Phys.* **356** (2017), no.3, 883–980.
- [12] J. Fröhlich, A. Knowles and S. Schwarz. On the mean-field limit of bosons Coulomb with two-body interaction. *Comm Math. Phys.* **288** (2009), 1023–1059.
- [13] J. Glimm and A Jaffe. Quantum Physics. A Functional Integral Point of View. *Springer-Verlag*, Second edition, 1987.
- [14] C. Kenig, G. Ponce, L. Vega. Well-posedness and scattering results for the generalized Korteweg-de Vries equation via contraction principle. *Comm. Pure Appl. Math.* **46** (1993), no. 4, 527–620.
- [15] C. Kenig, G. Ponce, L. Vega. The Cauchy problem for the Korteweg-de Vries equation in Sobolev spaces of negative indices. *Duke Math. J.* **71** (1993), no. 1, 1–21.
- [16] A. Knowles. Limiting dynamics in large quantum systems. ETH Zürich Doctoral Thesis (2009). ETHZ e-collection 18517.

- [17] J. Lebowitz, H. Rose, E. Speer. Statistical mechanics of the nonlinear Schrödinger equation. *J. Stat. Phys.* **50** (1988), 657–687.
- [18] M. Lewin, P. T. Nam, N. Rougerie. Derivation of nonlinear Gibbs measures from many-body quantum mechanics. *J. Éc. Polytech. Math* **2** (2015), 65–115.
- [19] M. Lewin, P. T. Nam, N. Rougerie. Gibbs measures based on 1D (an)harmonic oscillators as mean-field limits. *J. Math. Phys.* **59** (2018), no.4, 041901.
- [20] M. Lewin, P.T. Nam, N. Rougerie. Classical field theory limit of 2D many-body quantum Gibbs states. Preprint arXiv: 1805.08370.
- [21] H. P. McKean, K. L. Vaninsky. Action-angle variables for the cubic Schrödinger equation. *Comm. Pure Appl. Math* **50** (1997), no. 6, 489–562.
- [22] H. P. McKean, K. L. Vaninsky. Cubic Schrödinger: the petit canonical ensemble in action-angle variables. *Comm. Pure Appl. Math* **50** (1997), no. 7, 593–622.
- [23] M. Reed and B. Simon. Methods of modern mathematical physics, I: functional analysis. *Academic Press*, 1980.
- [24] B. Simon. The $P(\Phi)_2$ Euclidean (Quantum) Field Theory. *Princeton Univ. Press*, 1974.
- [25] B. Simon. Trace ideals and their applications. *Amer. Math. Soc.*, Second edition, 2005.
- [26] V. Sohinger. Bounds on the growth of high Sobolev norms of solutions to nonlinear Schrödinger equations. Ph.D. Thesis, MIT (2011).
- [27] T. Tao. Nonlinear dispersive equations: Local and global analysis. *AMS, CBMS Lecture Notes*, 2006.
- [28] P.E. Zhidkov. An invariant measure for the nonlinear Schrödinger equation. (Russian) *Dokl. Akad. Nauk SSSR* **317** (1991) 543–546; translation in *Soviet Math. Dokl.* **43**, 431–434.